

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/307718754>

Greenhouse gas emission factors associated with rewetting of organic soils

Article in *Mires and Peat* · April 2016

DOI: 10.19189/Map.2016.OMB.222

CITATIONS

154

READS

1,099

11 authors, including:



David Wilson

Earthy Matters Environmental Consultants

48 PUBLICATIONS 1,484 CITATIONS

[SEE PROFILE](#)



Dominique Blain

Environment and Climate Change Canada

12 PUBLICATIONS 751 CITATIONS

[SEE PROFILE](#)



Daniel Murdiyarso

Center for International Forestry Research Bogor Indonesia & Department of Geo...

261 PUBLICATIONS 16,052 CITATIONS

[SEE PROFILE](#)



Susan Page

University of Leicester

138 PUBLICATIONS 13,497 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Impact of Forest and Landscape Structure on Human Well-Being Experience - Mapping Forest Recreational Value [View project](#)



NEROS - Network Monitoring Rewetted and Restored Peatlands/Organic Soils for Climate and Biodiversity Benefits [View project](#)

Greenhouse gas emission factors associated with rewetting of organic soils

D. Wilson¹, D. Blain², J. Couwenberg³, C.D. Evans⁴, D. Murdiyarso^{5,6}, S.E. Page⁷,
F. Renou-Wilson⁸, J.O. Rieley⁹, A. Sirin¹⁰, M. Strack¹¹ and E.-S. Tuittila¹²

¹ Earthy Matters Environmental Consultants, Donegal, Ireland

² Environment Canada, Gatineau, Canada

³ University of Greifswald /Greifswald Mire Centre, Greifswald, Germany

⁴ Centre for Ecology and Hydrology, Bangor, Wales, UK

⁵ Center for International Forestry Research, Bogor, Indonesia

⁶ Bogor Agricultural University, Bogor, Indonesia

⁷ University of Leicester, Leicester, England, UK

⁸ University College Dublin, Dublin, Ireland

⁹ University of Nottingham, Nottingham, England, UK

¹⁰ Institute of Forest Science, Russian Academy of Sciences, Uspenskoe, Russia

¹¹ University of Waterloo, Ontario, Canada

¹² University of Eastern Finland, Joensuu, Finland

SUMMARY

Drained organic soils are a significant source of greenhouse gas (GHG) emissions to the atmosphere. Rewetting these soils may reduce GHG emissions and could also create suitable conditions for return of the carbon (C) sink function characteristic of undrained organic soils. In this article we expand on the work relating to rewetted organic soils that was carried out for the 2014 Intergovernmental Panel on Climate Change (IPCC) *Wetlands Supplement*. We describe the methods and scientific approach used to derive the Tier 1 emission factors (the rate of emission *per* unit of activity) for the full suite of GHG and waterborne C fluxes associated with rewetting of organic soils. We recorded a total of 352 GHG and waterborne annual flux data points from an extensive literature search and these were disaggregated by flux type (i.e. CO₂, CH₄, N₂O and DOC), climate zone and nutrient status. Our results showed fundamental differences between the GHG dynamics of drained and rewetted organic soils and, based on the 100 year global warming potential of each gas, indicated that rewetting of drained organic soils leads to: net annual removals of CO₂ in the majority of organic soil classes; an increase in annual CH₄ emissions; a decrease in N₂O and DOC losses; and a lowering of net GHG emissions. Data published since the *Wetlands Supplement* (n=58) generally support our derivations. Significant data gaps exist, particularly with regard to tropical organic soils, DOC and N₂O. We propose that the uncertainty associated with our derivations could be significantly reduced by the development of country specific emission factors that could in turn be disaggregated by factors such as vegetation composition, water table level, time since rewetting and previous land use history.

KEY WORDS: carbon, carbon dioxide, DOC, emission factors, methane, nitrous oxide, peat

INTRODUCTION

Organic soils store an estimated 600 Gt of carbon (C) worldwide (Page *et al.* 2011, Yu 2012), more than is currently held in the biomass of all the forests of the world (Köhl *et al.* 2015, Joosten *et al.* in press). Anthropogenic disturbance can destabilise these C stocks, often by accelerating decomposition of organic matter through drainage, which is commonly carried out to allow for conventional agricultural activities, forestry, infrastructure development or extraction of the peat.

Drained organic soils are significant sources of greenhouse gas (GHG) emissions to the atmosphere,

accounting for around 10 % of all GHG emissions from the agriculture, forestry and other land use (AFOLU) sectors (Smith *et al.* 2014). The decrease in water table levels following drainage leads to increased emissions of carbon dioxide (CO₂) (e.g. Chistotin *et al.* 2006, Maljanen *et al.* 2007, Jauhiainen *et al.* 2012, Salm *et al.* 2012, Renou-Wilson *et al.* 2014), methane (CH₄) ‘hotspots’ in drainage ditches (e.g. Minkinen & Laine 2006, Schrier-Uijl *et al.* 2011, Jauhiainen & Silvennoinen 2012, Sirin *et al.* 2012, Evans *et al.* 2015), reduced CH₄ emissions from drained land surfaces (e.g. Chistotin *et al.* 2006, Wilson *et al.* 2009) and high nitrous oxide (N₂O) emissions, particularly in

association with nutrient rich organic soils (e.g. Regina *et al.* 1996, van den Pol-van Dasselaar *et al.* 1998, Ernfors *et al.* 2008). In addition, drainage increases the vulnerability of organic soils to fire (Kettridge *et al.* 2015, Turetsky *et al.* 2015) which can lead to considerable additional GHG emissions, particularly from tropical organic soils (Page *et al.* 2002). Furthermore, waterborne C losses may be accentuated following drainage (Evans *et al.* 2015).

Rewetting of drained organic soils may reduce GHG emissions and waterborne C losses. Given the development of global climate policy and the high emissions associated with drained organic soils, it has been argued that rewetting and restoration of these soils should be included in mitigation strategies (Joosten *et al.* 2012, IPCC 2014a). Rewetting is the deliberate action of raising the water table in soils that have previously been drained for forestry, agriculture (crop production and grazing), water supply, peat extraction and other human-related activities, in order to re-establish and maintain water saturated conditions, e.g. by blocking drainage ditches, construction of bunds or disabling drainage pump facilities. Rewetting¹ can have several objectives such as nature conservation, GHG emission reductions and the promotion of leisure activities or paludiculture on saturated organic soils.

Under the United Nations Framework Convention on Climate Change (UNFCCC) and its Kyoto Protocol, Annex 1 countries prepare annual National Inventory Reports detailing GHG emissions and removals from six different sectors². While emissions from some drained organic soils are reported annually, the emissions and removals (i.e. C uptake) associated with rewetted organic soils have not been included thus far owing to an absence of data and a lack of methodological guidance (IPCC 2006). However, this gap has been addressed by the recent *2013 Supplement to the 2006 Guidelines for National Greenhouse Gas Inventories: Wetlands* (IPCC 2014a; hereafter referred to as the *Wetlands Supplement*), in which Tier 1 (default) emission factors are provided for

rewetted organic soils, disaggregated by climate region and nutrient status. The *Wetlands Supplement* provides methodological guidance for countries to report GHG emissions/removals arising from the rewetting of organic soils in their national inventory submissions. A second supplement (not discussed in this paper) provides guidance for accounting of these emissions/removals under Articles 3.3 and 3.4 of the Kyoto Protocol (second commitment period) (IPCC 2014b).

This article expands on the work relating to rewetted organic soils that was carried out for the *Wetlands Supplement*. Specifically, we describe here the methods and scientific approach used in the *Wetlands Supplement* to derive the Tier 1 emission factors for the full suite of GHG and waterborne C fluxes associated with rewetting of organic soils. We examine the robustness of the emission factors by comparing the Tier 1 values in the *Wetlands Supplement* with data published since that time. We compare GHG emissions and the global warming potential (GWP) of selected land use categories as defined by IPCC (2014a) under drained and rewetted conditions. Finally, we provide detailed information for countries that wish to move to higher reporting Tiers, identify current research gaps, and highlight priority areas for future research.

METHODS

In the land use sector, changes in C pools are commonly addressed using a stock-based approach: stocks are assessed at two points in time and the difference is interpreted as a flux of CO₂. However, C fluxes in organic soils are generally small relative to the total stock and are thus problematic to measure in a stock based approach. Net CO₂ emissions or removals from organic soils are more accurately measured directly as fluxes (IPCC 1996). Emissions (and removals) of the other main GHGs originating from the land use sector (CH₄ and N₂O) are commonly treated as fluxes. Fluxes are denoted here as CO₂-C, CH₄-C and N₂O-N. This notation is consistent with that used in the *2006 IPCC Guidelines for Greenhouse Gas Inventories* (IPCC 2006). We follow the sign convention that uses the atmosphere as a reference, whereby an emission is a positive flux from the soil to the atmosphere (and a removal is a negative flux).

The focus of this article is on GHG emissions/removals associated with *only* the soil C pool in rewetted organic soils. While the C contained in the woody biomass pool may be substantial in treed peatlands, particularly in the tropics, it is outside the scope of this paper. Instead,

¹ Wetland *restoration* always aims to permanently re-establish the pre-disturbance ecosystem, including the hydrological and biogeochemical processes typical of water saturated soils, as well as the vegetation cover that pre-dated the disturbance (Nelleman & Corcoran 2010). Normally, the restoration of drained organic soils is accompanied by rewetting, while the restoration of undrained but otherwise disturbed wetlands may not require rewetting. *Rehabilitation*, as defined by Poopathy *et al.* (2005) and Nelleman & Corcoran (2010), can involve a large variety of practices on formerly drained organic soils, which may or may not include rewetting. For example, the re-establishment of vegetation on a drained site without rewetting is a form of site rehabilitation.

² Energy; industrial processes; solvents and other product use; agriculture; land use, land use change and forestry (LULUCF); waste.

readers are directed to methodologies contained in the 2006 IPCC Guidelines (IPCC 2006) for quantifying C stock changes in the woody biomass and dead wood pools.

It is difficult to distinguish between the other C pools³ in organic soils. Living non-woody biomass (mosses, sedges, grasses) can be hard to separate from the dead litter derived from it, and the litter can be hard to separate from the (organic) soil. For example, the separation between the live and dead portions of mosses is not always clear; nor is it clear whether recently dead mosses should be included in the litter or the soil pool. Similarly, the (organic) soil in sedge dominated and forested tropical peatlands is made up mainly of dead root material and the distinction between the soil and recently dead roots is not easily made; nor is the distinction between recently dead and live roots straightforward. The default emission factors presented in this paper are all derived from published direct flux measurements (from eddy covariance (EC) towers and/or static chambers) over organic soils with moss and/or herbaceous and/or dwarf shrub vegetation. While some publications attempt to assess changes in the aboveground biomass pool separately from changes in the other pools, most combine all pools together. Therefore, we define the composite terms $CO_2-C_{composite}$ in [2], $CH_4-C_{composite}$ in [3] and $N_2O-N_{composite}$ in [4], which integrate all emissions (i.e. ecosystem respiration (autotrophic and heterotrophic), CH_4 production, denitrification, nitrification) and removals (i.e. photosynthesis, CH_4 oxidation) arising from the soil and the aboveground and belowground vegetation components other than trees.

Net annual C stock change

The net annual C stock change of rewetted organic soils ΔC_R (t C ha⁻¹ yr⁻¹) is the total net loss (or gain) of C from the soil (a loss is indicated by a positive value and a gain by a negative value) resulting from the balance between emissions and removals of both CO_2 and CH_4 , including on-site and off-site components:

$$\Delta C_R = CO_2-C_R + CH_4-C_R \quad [1]$$

where CO_2-C_R is the net flux of C as CO_2 (both on-site and off-site) from the rewetted organic soil (t C ha⁻¹ yr⁻¹) and CH_4-C_R is the net flux of CH_4 from the rewetted organic soil (kg C ha⁻¹ yr⁻¹).

³ The six pools in AFOLU are (1) aboveground biomass, (2) belowground biomass, (3) dead wood, (4) litter, (5) soil and (6) harvested wood products.

CO_2 emissions/removals

For carbon dioxide (CO_2) emissions/removals, we can write:

$$CO_2-C_R = CO_2-C_{composite} + CO_2-C_{DOC} + CO_2-C_{fire} \quad [2]$$

where $CO_2-C_{composite}$ denotes net CO_2 fluxes from the soil and non-tree vegetation (t C ha⁻¹ yr⁻¹), CO_2-C_{DOC} is off-site CO_2 emissions from dissolved organic carbon exported from rewetted organic soils (t C ha⁻¹ yr⁻¹) and CO_2-C_{fire} is C lost as CO_2 emissions from the burning of rewetted organic soils (t C ha⁻¹ yr⁻¹).

CH_4 emissions/removals

Methane (CH_4) emissions/removals from rewetted organic soils result from (a) the balance between biochemical CH_4 production and oxidation and (b) emissions of CH_4 produced by the combustion of soil organic matter during fire, and can be summarised by the equation

$$CH_4-C_R = CH_4-C_{composite} + CH_4-C_{fire} \quad [3]$$

where $CH_4-C_{composite}$ denotes net CH_4 fluxes from the soil and non-tree vegetation (kg C ha⁻¹ yr⁻¹), and CH_4-C_{fire} is C lost as CH_4 from the burning of rewetted organic soils (kg C ha⁻¹ yr⁻¹).

The CH_4 emission factors provided here relate to $CH_4-C_{composite}$ only (i.e. emissions from livestock and the burning of biomass are excluded). CH_4 emissions result from the decomposition of organic soil material by microbes under anoxic conditions, which is strongly controlled by the redox potential within the soil (Fiedler & Sommer 2000) and by soil temperature (van Winden *et al.* 2012). Emissions also originate from the partial decay of non-tree vegetation, but since this cannot easily be separated from the organic soil the emissions are combined here as $CH_4-C_{composite}$. The probability of fire in rewetted organic soils is low because the water table should be near the surface, but possible soil emissions from fires are included in [3] for completeness. High spatial variation in microtopography, water table depth and vegetation cover is typical of undrained organic soils and is reflected in CH_4 fluxes (Strack *et al.* 2006, A. Laine *et al.* 2007a, Riutta *et al.* 2007, Maanaviija *et al.* 2011). Rewetting recreates this natural heterogeneity to some extent, with blocked ditches forming the wetter end of the variation (Strack & Zuback 2013). For this reason, former ditches are included as a part of rewetted sites and not treated separately.

N₂O emissions/removals

The emissions of N₂O from rewetted organic soils are controlled by the quantity of nitrogen available for nitrification and denitrification, and the redox potential. They are summarised by:

$$N_2O-N_R = N_2O-N_{composite} + N_2O-N_{fire} \quad [4]$$

where N_2O-N_R denotes the net flux of N as N₂O from rewetted organic soils (kg N ha⁻¹ yr⁻¹), $N_2O-N_{composite}$ is net N₂O fluxes from the soil and non-tree vegetation (kg N ha⁻¹ yr⁻¹), and N_2O-N_{fire} is N lost as N₂O from the burning of rewetted organic soils (kg N ha⁻¹ yr⁻¹). Published data are currently insufficient to develop default N₂O emission factors for the burning of organic soils (see Chapter 2, IPCC 2014a). Therefore, N_2O-N_{fire} is not considered further here.

Derivation of emission factors

An extensive literature review was conducted to collate all GHG (CO₂, CH₄, N₂O) and DOC studies that were available at the time the *Wetlands Supplement* was prepared (i.e. 2013) for (1) rewetted organic soils (includes rewetted, restored and wet managed sites) and (2) natural/undrained organic soils (to be used as proxies for rewetted soils, see criteria below). Laboratory and manipulation experiments were excluded. Literature sources included both peer reviewed and ‘grey literature’ (i.e. not peer reviewed) studies. In the case of the latter, we reviewed the studies and expert judgement was exercised as to whether they were scientifically acceptable for inclusion. In total, three grey literature studies were included.

No studies exist on rewetted tropical sites with the water table close to the surface. For temperate and boreal sites we plotted annual CO₂ and CH₄ fluxes against the mean water table level (*MWTL*) (positive values upwards from the soil surface) for both natural/undrained soils and rewetted soils to assess whether natural/undrained organic soils could function as proxies for rewetted organic soils. In temperate regions *MWTL* was calculated over one year where the flux measurements covered the full 12 months, while in boreal regions the *MWTL* applied to the growing season only. The non-normally distributed CH₄ fluxes (assessed using the Kolmogorov-Smirnov test) were log transformed prior to regression analyses. Fitted linear regression lines (CO₂ and log₁₀CH₄ fluxes = $a + (b \times MWTL)$) were compared for each climate zone. Differences between the undrained and rewetted groups were compared using General Linear Models (IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY,

USA) to examine the homogeneity of the regression slopes. The 95 % confidence intervals of the intercepts (parameter a) were also compared.

All studies included in the database reported GHG flux estimates using either static chamber or EC techniques. The chamber method involves the measurement of gas fluxes at high spatial resolution and is widely employed in circumstances where vegetation is either short (i.e. of low stature) or absent. EC towers are typically used at sites that are relatively flat and homogeneous, which includes open and treed organic soils. For a more detailed description of both methodologies see Alm *et al.* (2007).

A detailed database of annual GHG fluxes was then constructed to evaluate the main drivers of GHG dynamics in rewetted organic soils. When available, the following attributes were extracted from the literature source and included in the database for analysis: climate zone as defined by IPCC (2006), nutrient status⁴, mean and median water table level (as well as minimum and maximum values), soil pH, thickness of the organic soil layer, C/N quotient, degree of humification, soil moisture, soil bulk density, plant cover and species or functional groups, previous land use and time since rewetting. The criteria for inclusion in the database were as follows:

- (1) The study reported GHG fluxes from rewetted organic soils, abandoned and spontaneously rewetted organic soils, and natural (undrained) organic soils. Natural sites with *MWTL* lower than -30 cm were designated as ‘not wet’, and were not included in the final database. In other words, only natural sites with an annual or seasonal *MWTL* of -30 cm or shallower (i.e. closer to or above the soil surface) were deemed suitable as proxies for rewetted sites since the *MWTL* recorded at all the rewetted sites in our database was at or shallower than -30 cm. Studies from rewetted sites with *MWTL* more than 10 cm above the soil surface were not included either, as they were judged to refer to flooded land. Flooded land is considered a separate type of land in IPCC terms (IPCC 2006).
- (2) The study had to report either seasonal or annual GHG fluxes. Studies in the database that reported

⁴ Nutrient poor organic soils (bog) predominate in boreal regions, while in temperate regions nutrient rich (fen) sites are more common. In many cases, nutrient poor organic soil layers are underlain by nutrient rich layers; in some situations, after industrial extraction of the nutrient poor top layers the rewetted residual soil layers may be considered nutrient rich due to the influence of groundwater and the high nutrient status of the bottom layers.

daily flux values were not used, as upscaling to annual flux values could have led to very high errors. However, non-annualised CH₄ flux values were used for tropical sites as annual data from those sites are scarce and seasonality is either absent or relates to wet and dry seasons only. During the dry season some tropical sites show very large (> 1 m) drops in water table level. We discarded the measurements made during these times because the conditions cannot be deemed wet, and accepted that the omission of data for naturally dry periods may result in slight overestimation of the CH₄ emission factor. In boreal sites, seasonal CH₄ fluxes (typically May to October) were converted to annual fluxes by assuming that an additional 15 % of flux occurs in the non-growing season (Saarnio *et al.* 2007). For CO₂, seasonal flux data were converted to annual fluxes by adding 15 % to the seasonal ecosystem respiration data from each study (Saarnio *et al.* 2007) and making the assumption that no photosynthesis occurs under snow. This adjustment may result in a slight overestimation of losses outwith the growing season because photosynthesis (and hence C uptake) may occur for a short time outside the period covered by seasonal studies. For studies where ecosystem respiration data were not explicitly reported, a value of 0.30 t CO₂-C ha⁻¹ (30g CO₂-C m⁻²) for non-growing-season respiration was used (Saarnio *et al.* 2007).

- (3) Studies had to indicate a *MWTL* for each annual GHG flux reported. In some cases, the GHG flux value was accepted for inclusion on the basis that water table information was available from other publications.
- (4) For studies using the EC technique, care was taken not to use annual CO₂ fluxes that included accumulation of C into the tree biomass pool (e.g. in the case of treed organic soils) as this would have resulted in double counting under IPCC rules. As such, our derived default emission factors for CO₂ exclude CO₂ uptake into the tree biomass.
- (5) Rewetting as a management practice is in its infancy in the tropics, and while projects to rehabilitate drained peatlands are being initiated in south-east Asia, there are no published flux measurement data for successfully rewetted tropical organic soils from which to derive emission factors. Therefore, a default emission factor for rewetted tropical organic soils was developed based on surrogate data. Subsidence measurements provide a good measure of C

losses from drained organic soils (see Chapter 2 in IPCC 2014a) and in tropical organic soils subsidence is near zero when the water table approaches the surface (Hooijer *et al.* 2012, see also Couwenberg *et al.* 2010). In light of the available evidence, the Tier 1 default emission factor was set at 0 t CO₂-C ha⁻¹ yr⁻¹. This value is consistent with observations of subsidence and reflects the fact that rewetting effectively stops soil organic matter oxidation but does not necessarily re-establish the soil C sink function.

Data treatment

In the database, flux values were standardised to the following units; t CO₂-C ha⁻¹ yr⁻¹, kg CH₄-C ha⁻¹ yr⁻¹, t C ha⁻¹ yr⁻¹ (for DOC) and kg N₂O-N ha⁻¹ yr⁻¹. For multi-year studies from the same site, annual flux estimates were averaged over the years. Emission factors were calculated as mean fluxes, with 95 % confidence intervals calculated for each of the categories. In order to examine the robustness of the derived emission factor values, we conducted a similar literature search for publications that might have been missed in the original literature search and for new datasets published since the *Wetlands Supplement*.

Comparison with drained organic soils

To assess the impact of rewetting on GHG dynamics in organic soils, the emission factors derived in Chapter 2 of the *Wetlands Supplement* (IPCC 2014a) for drained land use categories were compared to the emission factors derived here for their rewetted counterparts. The net GHG emissions, based on the global warming potential (GWP; t CO₂-eq ha⁻¹ yr⁻¹) of each gas, were calculated for the land use categories under drained and rewetted conditions. CH₄ and N₂O fluxes were converted to CO₂ equivalents according to their GWP on a 100-year timescale including climate-carbon feedbacks: CH₄ = 34 and N₂O = 298 (Myhre *et al.* 2013).

RESULTS

Undrained sites as proxies

The relationships between *MWTL* and CO₂/CH₄ fluxes were very similar for undrained and rewetted sites in the boreal and temperate climate zones (Figure 1, Figure 2). In the boreal zone, the change in magnitude of the CO₂ flux *per* unit change in *MWTL* was small (Figure 1a), whereas it was more pronounced in the temperate zone (Figure 1b). Variance was clearly larger in the temperate data than in the boreal data. CH₄ fluxes in both climate




Figure 1. Relationship between annual carbon dioxide (CO_2) flux ($\text{t C ha}^{-1} \text{ yr}^{-1}$) and mean water table level ($MWTL$, cm) for undrained and rewetted organic soils in (a) boreal and (b) temperate climate zones. Solid grey and dashed lines are the regression lines for the undrained and rewetted sites respectively. Regression equations and parameters are shown in Table 1. Negative $MWTL$ indicates mean water table level below the soil surface and positive $MWTL$ indicates mean water table above the soil surface. Negative CO_2 values indicate annual CO_2 removals and positive values indicate annual CO_2 emissions.




Figure 2. Relationships between annual methane (CH_4) fluxes ($\log_{10}(1 + (\text{kg C ha}^{-1} \text{ yr}^{-1})))$ and mean water table level ($MWTL$, cm) for undrained and rewetted organic soils in (a) boreal and (b) temperate climate zones. Solid grey and dashed lines are the regression lines for the undrained and rewetted sites respectively. Regression equations and parameters are shown in Table 1. Negative $MWTL$ indicates mean water table level below the soil surface and positive $MWTL$ indicates mean water table above the soil surface. Positive CH_4 values indicate annual emissions and negative values indicate annual removals.

zones showed strong sensitivity to changes in $MWTL$, with lower fluxes observed in conjunction with deeper water table levels (Figure 2). No significant difference in the homogeneity of regression slopes was observed (Table 1) and the 95 % confidence intervals of the intercepts (parameter a) overlapped in comparable datasets; so we concluded that undrained sites can, for the purposes of emission factor calculations, act as

proxies for rewetted sites with the same $MWTL$, and thereafter we combined the two datasets. Moreover, given the observed similarity between rewetted and undrained soils in the boreal and temperate climate zones, in the absence of existing data we also made the highly generalised assumption that, for the derivation of CH_4 and N_2O emission factors, tropical undrained organic soils can act as proxies for tropical rewetted organic soils.

Table 1. Results of General Linear Models (model equations: $\text{flux} = a + (b \times \text{MWTL})$) used to test for homogeneity of regression slopes between undrained and rewetted sites in the boreal and temperate climate zones. Test results are given as F and p values. Standard errors of the model parameters are shown in parentheses. Note: sample size (n) includes all annual flux measurements (i.e. not averaged multi-year datasets from the same site) recorded in the *Wetlands Supplement* database and, therefore, differs from the sample sizes described in Figure 3.

		Sample size	Model parameters		F	p
		n	a	b		
CO_2			(t C ha ⁻¹ yr ⁻¹)			
Boreal	Undrained	82	-0.61 (0.10)	-0.02 (0.01)	0.02	0.90
	Rewetted	26	-0.58 (0.18)	-0.02 (0.02)		
Temperate	Undrained	52	-0.29 (0.28)	-0.05 (0.02)	0.03	0.86
	Rewetted	64	-0.43 (0.29)	-0.04 (0.02)		
CH_4			(log ₁₀ (1+(kg C ha ⁻¹ yr ⁻¹)))			
Boreal	Undrained	49	2.27 (0.11)	0.052 (0.007)	0.01	0.92
	Rewetted	15	1.70 (0.45)	0.049 (0.033)		
Temperate	Undrained	47	1.99 (0.15)	0.043 (0.016)	0.10	0.76
	Rewetted	38	1.88 (0.14)	0.036 (0.012)		

Data distribution

A total of 123 (CO_2), 164 (CH_4), 36 (N_2O) and 29 (DOC) data entries satisfied the criteria outlined above and were included in the final database (Figure 3). Data entries for CO_2 and CH_4 were relatively evenly spread between the boreal and temperate zones, while no CO_2 entries and only a small number (11) of CH_4 entries were recorded for the tropical zone. In the *Wetlands Supplement*, only two N_2O entries were reported for rewetted organic soils (Hendriks *et al.* 2007, Wilson *et al.* 2013), and we provide here a further 32 data entries mainly from the temperate and boreal zones. DOC entries were found for all three climate zones. In most cases (with the exceptions of tropical data and DOC entries), the data could be further disaggregated by nutrient status (i.e. nutrient poor (bog), nutrient rich (fen)) (Figure 3). Largely due to the small number of data points within any given category, there was insufficient evidence to support the disaggregation of data by additional site conditions, previous land use, time since rewetting or drainage status (i.e. undrained and rewetted).

The CO_2 flux data were normally distributed (Shapiro-Wilk test; $p > 0.05$) when disaggregated by climate zone (Figure 4a). The range of values was markedly wider in the temperate zone data (-4.4 to 4.78 t C ha⁻¹ yr⁻¹) than in the boreal data (-2.68 to 1.48 t C ha⁻¹ yr⁻¹). The CH_4 flux data for both boreal and temperate zones had a log normal distribution characterised by a high number of flux values close



Figure 3. Number of data entries (see main text for inclusion criteria) used to derive carbon dioxide (CO_2), methane (CH_4), nitrous oxide (N_2O) and dissolved organic carbon (DOC) emission factors. Entries are disaggregated by climate zone (boreal, temperate and tropical), nutrient status (NP = nutrient poor, NR = nutrient rich) and drainage status (U = undrained, R = rewetted).




Figure 4. Frequency distributions of (a) carbon dioxide (CO_2 ; $\text{t C ha}^{-1} \text{ yr}^{-1}$), (b) methane (CH_4 ; $\text{kg C ha}^{-1} \text{ yr}^{-1}$), (c) dissolved organic carbon (DOC; $\text{t C ha}^{-1} \text{ yr}^{-1}$) and (d) nitrous oxide (N_2O ; $\text{kg N ha}^{-1} \text{ yr}^{-1}$) fluxes in undrained and rewetted organic soils in the boreal, temperate and tropical (DOC and N_2O) climate zones. Negative values indicate annual removals and positive values indicate annual emissions.

to zero together with some very high values (Figure 4b). As with the CO_2 data, the range of CH_4 flux values was larger in the temperate zone than in the boreal (Figure 4b). DOC data were normally distributed (Shapiro-Wilk test; $p > 0.05$) in all three climate zones (Figure 4c), with the widest range of values observed in the temperate zone data (0.05 to $0.61 \text{ t C ha}^{-1} \text{ yr}^{-1}$). Similarly, N_2O flux values were mainly congregated around zero with the exception of one high value from a sago (*Metroxylon sagu*) plantation in the tropical zone (Figure 4d).

Emission factors

CO_2 - $C_{\text{composite}}$

The CO_2 - $C_{\text{composite}}$ values varied considerably across climate zones and nutrient status (Table 2). Boreal nutrient rich sites showed the highest annual CO_2 removals at $0.55 \text{ t C ha}^{-1} \text{ yr}^{-1}$ and temperate nutrient rich sites showed the highest annual emissions of $0.50 \text{ t C ha}^{-1} \text{ yr}^{-1}$. Boreal and temperate nutrient poor sites showed CO_2 removal values of 0.34 and $0.24 \text{ t C ha}^{-1} \text{ yr}^{-1}$ respectively. Uncertainty associated with the CO_2 - $C_{\text{composite}}$ values was lowest for boreal nutrient rich sites ($\pm 40\%$) and highest in the temperate nutrient rich sites ($\pm 242\%$).

CH_4 - $C_{\text{composite}}$

The lowest emissions were observed in the boreal nutrient poor and tropical sites ($41 \text{ kg C ha}^{-1} \text{ yr}^{-1}$) and the highest emissions were seen in the temperate nutrient rich sites ($216 \text{ kg C ha}^{-1} \text{ yr}^{-1}$). Associated uncertainty was very high across all groups (Table 2).

CO_2 - C_{DOC}

The DOC data did not support disaggregation by nutrient status. DOC flux values were 0.08 , 0.26 and $0.57 \text{ t C ha}^{-1} \text{ yr}^{-1}$ from the boreal, temperate and tropical zones, respectively (Table 3). The parameter $\text{Frac}_{\text{DOC-CO}_2}$ sets the proportion of DOC exported from organic soils that is ultimately emitted as CO_2 . A value of zero would coincide with all the DOC export being deposited in stable forms in lake or marine sediments; as this would simply represent a translocation of C between stable stores, it would not need to be estimated. However, most data on DOC processing indicate that a high proportion is converted to CO_2 in headwaters, rivers, lakes and coastal seas (IPCC 2014a). A value of 0.9 is proposed for $\text{Frac}_{\text{DOC-CO}_2}$ with an uncertainty range of 0.8 to 1 (IPCC 2014a, Evans *et al.* 2015). This resulted in CO_2 - C_{DOC} values of 0.08 , 0.24 and $0.51 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for the boreal, temperate and tropical zones, respectively (Table 3).

N_2O - $N_{\text{composite}}$

As the sample sizes were low, it was not possible to derive a robust N_2O - $N_{\text{composite}}$ value disaggregated by nutrient status (Table 4). Values for the boreal and temperate zones were very similar (0.06 – $0.07 \text{ kg N ha}^{-1} \text{ yr}^{-1}$). Average emissions from the tropical zone were higher at $0.94 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ but displayed very high uncertainty due to the very small sample size ($n = 5$) and the inclusion of a single high data point (see Melling *et al.* 2007).

Table 2. Emission factors for carbon dioxide ($CO_2-C_{composite}$: t C ha⁻¹ yr⁻¹) and methane ($CH_4-C_{composite}$: kg C ha⁻¹ yr⁻¹) in wet organic soils, and associated uncertainty ranges (95 % confidence intervals). Emission factors disaggregated by climate zone and by nutrient status where appropriate. Positive emission factors indicate emissions to the atmosphere and negative values indicate removals from the atmosphere.

Climate zone	Nutrient status	$CO_2-C_{composite}$ (t C ha ⁻¹ yr ⁻¹)	95 % range	$CH_4-C_{composite}$ (kg C ha ⁻¹ yr ⁻¹)	95 % range
Boreal*	Poor	-0.34	-0.59 to -0.09	41	0.5 to 246
	Rich	-0.55	-0.77 to -0.34	137	0 to 493
Temperate**	Poor	-0.23	-0.64 to +0.18	92	3 to 445
	Rich	0.50	-0.71 to +1.71	216	0 to 856
Tropical***		0		41	7 to 134

* $CO_2-C_{composite}$ derived from Alm *et al.* 1997, J. Laine *et al.* 1997, Suyker *et al.* 1997, Bubier *et al.* 1999, Komulainen *et al.* 1999, Soegaard & Nordstroem 1999, Tuittila *et al.* 1999, Waddington & Price 2000, Waddington & Roulet 2000, Whiting & Chanton 2001, Heikkinen *et al.* 2002, Harazono *et al.* 2003, Nykänen *et al.* 2003, Yli-Petäys *et al.* 2007, Kivimäki *et al.* 2008, Nilsson *et al.* 2008, Sagerfors *et al.* 2008, Aurela *et al.* 2009, Drewer *et al.* 2010, Soini *et al.* 2010, Maanavilja *et al.* 2011.

$CH_4-C_{composite}$ derived from the following source material Clymo & Reddaway 1971, Verma *et al.* 1992, Bubier *et al.* 1993, Nykänen *et al.* 1995, J. Laine *et al.* 1996, Alm *et al.* 1997, Komulainen *et al.* 1998, Tuittila *et al.* 2000, Waddington & Roulet 2000, Whiting & Chanton 2001, Gauci *et al.* 2002, Yli-Petäys *et al.* 2007, Drewer *et al.* 2010, Juottonen *et al.* 2012, Urbanová 2012, Strack & Zuback 2013.

** $CO_2-C_{composite}$ derived from Shurpali *et al.* 1995, Lafleur *et al.* 2001, Wickland 2001, Aurela *et al.* 2002, Schulze *et al.* 2002, Petrone *et al.* 2003, Roehm & Roulet 2003, Billett *et al.* 2004, Drösler 2005, Nagata *et al.* 2005, Bortoluzzi *et al.* 2006, Hendriks *et al.* 2007, Jacobs *et al.* 2007, Lund *et al.* 2007, Riutta *et al.* 2007, Roulet *et al.* 2007, Wilson *et al.* 2007, Cagampan & Waddington 2008, Golovatskaya & Dyukarev 2009, Kurbatova *et al.* 2009, Drewer *et al.* 2010, Waddington *et al.* 2010, Adkinson *et al.* 2011, Couwenberg *et al.* 2011, Koehler *et al.* 2011, Christensen *et al.* 2012, Urbanová 2012, Drösler *et al.* 2013, Herbst *et al.* 2013, Strack & Zuback 2013, Wilson *et al.* 2013.

$CH_4-C_{composite}$ derived from Bartlett & Harriss 1993, Dise *et al.* 1993, Shannon & White 1994, Flessa *et al.* 1997, Augustin & Merbach 1998, Waddington & Price 2000, Wickland 2001, Wild *et al.* 2001, Augustin 2003, Sommer *et al.* 2004, von Arnold 2004, Cleary *et al.* 2005, Drösler 2005, Nagata *et al.* 2005, Augustin *et al.* 2006, Bortoluzzi *et al.* 2006, Hendriks *et al.* 2007, Jungkunst & Fiedler 2007, Roulet *et al.* 2007, Scottish Executive 2007, Nilsson *et al.* 2008, Tauchnitz *et al.* 2008, Wilson *et al.* 2009, Couwenberg *et al.* 2011, Glatzel *et al.* 2011, Koehler *et al.* 2011, Beetz *et al.* 2013, Drösler *et al.* 2013, Wilson *et al.* 2013.

*** For tropical rewetted organic soils where decayed organic material is not oxidised due to saturated conditions. $CH_4-C_{composite}$ derived from Inubushi *et al.* 1998, Hadi *et al.* 2001, Jauhiainen *et al.* 2001, Jauhiainen *et al.* 2004, Furukawa *et al.* 2005, Hadi *et al.* 2005, Jauhiainen *et al.* 2005, Jauhiainen *et al.* 2008, Melling *et al.* 2012, Pangala *et al.* 2013.

Table 3. Emission factors for dissolved organic carbon (DOC : t C ha⁻¹ yr⁻¹) and CO_2-C_{DOC} (t CO₂-C ha⁻¹ yr⁻¹) and associated uncertainty ranges (95 % confidence intervals). Note rounding artefact on DOC values for boreal climate zone.

Climate zone	DOC (t C ha ⁻¹ yr ⁻¹)	95 % range	CO_2-C_{DOC} (t CO ₂ -C ha ⁻¹ yr ⁻¹)	95 % range
Boreal*	0.08	0.06 to 0.11	0.08	0.05–0.11
Temperate**	0.26	0.17 to 0.36	0.24	0.14–0.36
Tropical***	0.57	0.49 to 0.64	0.51	0.40–0.64

* Derived from Koprivnjak & Moore 1992, Moore *et al.* 2003, Kortelainen *et al.* 2006, Ågren *et al.* 2008, Nilsson *et al.* 2008, Jager *et al.* 2009, Rantakari *et al.* 2010, Juutinen *et al.* 2013.

** Derived from Urban *et al.* 1989, Kolka *et al.* 1999, Clair *et al.* 2002, Moore *et al.* 2003, Dawson *et al.* 2004, Roulet *et al.* 2007, O'Brien 2009, Strack *et al.* 2008, Waddington *et al.* 2008, Koehler *et al.* 2009, Billet *et al.* 2010, di Folco & Kirkpatrick 2011, Dinsmore *et al.* 2011, Koehler *et al.* 2011, Strack & Zuback 2013, Turner *et al.* 2013.

*** Derived from Zulkifli 2002, Alkhatib *et al.* 2007, Baum *et al.* 2007, Yule & Gomez 2009, Moore *et al.* 2013.

Table 4. Emission factors for nitrous oxide (N_2O - $N_{composite}$; $kg\ N\ ha^{-1}\ yr^{-1}$) in rewetted organic soils and associated uncertainty range (95 % confidence intervals).

Climate zone	N_2O - $N_{composite}$	95 % range ($kg\ N\ ha^{-1}\ yr^{-1}$)
Boreal*	0.06	0.00 to 0.14
Temperate**	0.07	-0.03 to 0.14
Tropical***	0.94	-0.03 to 1.90

*Derived from Martikainen *et al.* 1995, Nykänen *et al.* 1995, J. Laine *et al.* 1996, Drewer *et al.* 2010, Salm *et al.* 2012.

**Derived from Wild *et al.* 2001, Drösler 2005, Nagata *et al.* 2005, Hendriks *et al.* 2007, Tauchnitz *et al.* 2008, Drewer *et al.* 2010, Beetz *et al.* 2013, Juszczak & Augustin 2013, Wilson *et al.* 2013, Beyer & Höper 2015, Günther *et al.* 2015, Minke *et al.* 2015, Vanselow-Algan *et al.* 2015.

***Derived from Inubushi *et al.* 2003, Melling *et al.* 2007, D. Murdiyarto unpublished data.

Robustness of EF derivation

We examined the robustness of the CO_2 and CH_4 emission factors by comparing the values derived above for the *Wetlands Supplement* with data published since that time (Figure 5). The variance in the new CO_2 and CH_4 values was high, but lay within that of the original dataset. New values for undrained and rewetted sites did not differ significantly from the respective original datasets (Figure 5). The majority of the new CO_2 data values were negative (i.e. CO_2 removal). Whereas the original dataset provided a positive (i.e. CO_2 emission) CO_2 - $C_{composite}$ value for temperate nutrient rich sites, the new data indicate that this may have been an overestimation of CO_2 emissions from these sites (Figures 5a and 6). For both CO_2 and CH_4 , the updated mean flux values (calculated from both the *Wetlands Supplement* data and the new data) for each of the five land types was within the 95 % confidence intervals of the original emission factors (Table 5).

Comparison with drained organic soils

CO_2 emissions decreased considerably following rewetting of drained organic soils (Figure 6a and Table 5). In contrast, CH_4 emissions were much higher following rewetting (Figure 6b and Table 5) and were also characterised by very high variability (Figure 6b). N_2O emissions were considerably reduced following rewetting (Figure 6c). The soil GWP was largely dominated by CH_4 emissions in the rewetted sites, and by CO_2 emissions in the




Figure 5. (a) Carbon dioxide (CO_2 ; $t\ C\ ha^{-1}\ yr^{-1}$) and (b) methane (CH_4 ; $kg\ C\ ha^{-1}\ yr^{-1}$) fluxes disaggregated by climate zone and nutrient status (NP = nutrient poor, NR = nutrient rich). Negative values indicate annual removals and positive values indicate annual emissions. *Wetlands Supplement* data are shown as grey circles and new data as red circles. Black horizontal line indicates the mean value (derived emission factor) calculated for the *Wetlands Supplement* data.

For literature used in the derivation of emission factors for the *Wetlands Supplement* (grey circles), see Tables 2 and 3. **New CO_2 values** taken from Neuzil 1997, Page *et al.* 2004, Chimner & Karberg 2008, Lähteenoja *et al.* 2009, Dommain *et al.* 2011, Lähteenoja *et al.* 2012, Lund *et al.* 2012, Beetz *et al.* 2013, Gažovič *et al.* 2013, Koebsch *et al.* 2013a, Olson *et al.* 2013, Urbanová *et al.* 2013b, Campbell *et al.* 2014, Cliche Trudeau *et al.* 2014, Humphreys *et al.* 2014, McVeigh *et al.* 2014, Peichl *et al.* 2014, Strack *et al.* 2014, Aurela *et al.* 2015, Dommain *et al.* 2015, Günther *et al.* 2015, Helfter *et al.* 2015, Hribljan *et al.* 2015, Kareksela *et al.* 2015, Levy & Gray 2015, Lund *et al.* 2015, Minke *et al.* 2015, Vanselow-Algan *et al.* 2015, C.D. Evans *et al.* in preparation, D. Wilson unpublished data. **New CH_4 values** taken from Inubushi *et al.* 2003, Melling *et al.* 2005, Rinne *et al.* 2007, Hirano *et al.* 2009, Long *et al.* 2010, Forbrich *et al.* 2011, Cliche Trudeau *et al.* 2013, Gažovič *et al.* 2013, Huth *et al.* 2013, Juszczak & Augustin 2013, Koebsch *et al.* 2013b, Nadeau *et al.* 2013, Olson *et al.* 2013, Urbanová *et al.* 2013a, Urbanová *et al.* 2013b, Adji *et al.* 2014, Ballantyne *et al.* 2014, Cooper *et al.* 2014, Günther *et al.* 2014, Strack *et al.* 2014, Beyer & Höper 2015, Goodrich *et al.* 2015, Koebsch *et al.* 2015, Levy & Gray 2015, Minke *et al.* 2015.

drained sites, although N_2O emissions were also significant in the latter (Table 5). Both the drained and rewetted sites had a net warming effect on the

Table 5. Global warming potential (GWP) for drained and rewetted (presented here using the derived value from the *Wetlands Supplement* and an updated value that incorporates the new data published post 2013) organic soils for selected land use categories as defined by IPCC (2014a). Methane (CH₄) fluxes include emissions from ditches and were calculated using the default ditch area provided by IPCC (2014a). CH₄ and nitrous oxide (N₂O) fluxes are converted to CO₂ eq. (t CO₂-eq ha⁻¹ yr⁻¹) according to their GWPs on a 100-year timescale including climate–carbon feedbacks: CH₄ = 34 and N₂O = 298 (Myhre *et al.* 2013). Positive values indicate a net warming effect on the climate and negative values indicate a net cooling effect. ER=emission reduction, NR=nutrient rich, NP=nutrient poor, DD=deeply drained, SD=shallow drained. Values in bold indicate the dominant GHG within each land use category.

	Land use category	Drained					Rewetted						Rewetted (updated)					
		CO ₂	DOC	CH ₄	N ₂ O	GWP	CO ₂	DOC	CH ₄	N ₂ O	GWP	ER	CO ₂	DOC	CH ₄	N ₂ O	GWP	ER
boreal	Forest Land NP	0.92	0.44	0.42	0.10	1.88	-1.25	0.29	1.86	0	0.90	0.98	-1.52	0.29	1.87	0.03	0.67	1.21
	Forest Land NR	3.41	0.44	0.25	1.50	5.60	-2.02	0.29	6.21	0	4.48	1.12	-1.93	0.29	5.64	0.03	4.03	1.57
	Cropland	28.97	0.44	1.98	6.09	37.48	-2.02	0.29	6.21	0	4.48	33.00	-1.93	0.29	5.64	0.03	4.03	33.45
	Grassland	20.90	0.44	2.03	4.45	27.82	-2.02	0.29	6.21	0	4.48	23.34	-1.93	0.29	5.64	0.03	4.03	23.79
	Peat extraction	10.27	0.44	1.12	0.14	11.97	-1.25	0.29	1.86	0	0.91	11.06	-1.52	0.29	1.87	0.03	0.67	11.30
temperate	Forest Land NP	9.53	1.14	0.27	1.31	12.25	-0.84	0.88	4.17	0	4.21	8.04	-1.22	0.88	4.09	0.03	3.78	8.47
	Forest Land NR	9.53	1.14	0.27	1.31	12.25	1.83	0.88	9.79	0	12.50	-0.25	0.96	0.88	10.7	0.03	12.57	-0.32
	Cropland	28.97	1.14	1.98	6.09	38.18	1.83	0.88	9.79	0	12.50	25.68	0.96	0.88	10.7	0.03	12.57	25.61
	Grassland NP	19.43	1.14	2.04	2.01	24.62	-0.84	0.88	4.17	0	4.21	20.41	-1.22	0.88	4.09	0.03	3.78	20.84
	Grassland NR, DD	22.37	1.14	2.50	3.84	29.85	1.83	0.88	9.79	0	12.50	17.35	0.96	0.88	10.7	0.03	12.57	17.28
	Grassland NR, SD	13.20	1.14	2.16	0.75	17.25	1.83	0.88	9.79	0	12.50	4.75	0.96	0.88	10.7	0.03	12.57	4.68
	Peat extraction	10.27	1.14	1.12	0.14	12.67	-0.84	0.88	4.17	0	4.21	8.46	-1.22	0.88	4.09	0.03	3.78	8.89
tropical	Plantation	55.00	3.01	1.58	0.56	60.15	0.00	2.09	1.86	0	3.95	56.20	0.00	2.09	2.77	0.44	5.30	54.85
	Cropland	51.33	3.01	1.77	2.34	58.45	0.00	2.09	1.86	0	3.95	54.50	0.00	2.09	2.77	0.44	5.30	53.15

Note that CO₂ and DOC values for Rewetted (updated) are the same as Rewetted for tropical land use categories, and that GHG and GWP values are for soil emissions/removals and do not take into account C emissions/removals associated with the woody biomass pool.




Figure 6. Mean annual (a) carbon dioxide (CO₂; t C ha⁻¹ yr⁻¹), (b) methane (CH₄; kg C ha⁻¹ yr⁻¹) and (c) nitrous oxide (N₂O; kg N ha⁻¹ yr⁻¹) emission factors for drained (values taken from IPCC 2014a, Chapter 2) and rewetted (presented using the derived value from the *Wetlands Supplement* and an updated value that incorporates the new data published post 2013) organic soils for selected land use categories as defined by IPCC (2014a). Error bars represent the 95 % confidence intervals. Positive values indicate annual emissions and negative values indicate annual removals. NR = nutrient rich, NP = nutrient poor.

climate, although rewetting did result in large emissions reductions across most land use categories (Table 5). However, higher emissions were estimated from the rewetted temperate nutrient rich forest sites compared to their drained counterparts, an apparent anomaly caused by the lack of disaggregation by nutrient status in the drained sites (due to a low sample size), compounded by disaggregation in the rewetted sites.

DISCUSSION

With the exception of temperate nutrient rich organic soils, all rewetted organic soils were estimated to be net CO₂ sinks, although the inclusion of the most recent data suggests that CO₂ emissions from temperate nutrient rich soils may be much lower than those derived from the original dataset, with many studies indicating a net CO₂ sink (Figure 5a). Since undrained temperate nutrient rich organic soils must (in order to exist) have acted as net CO₂ sinks during the course of their formation, it seems reasonable to assume that a successfully rewetted nutrient rich peatland should also ultimately act as a CO₂ sink, or at least cease to act as a CO₂ source. The positive emission factor associated with this category may therefore be a consequence of incorporating data from studies

undertaken on incompletely rewetted sites, sites at which the natural peat-forming vegetation has not re-established, or recently rewetted sites that are still undergoing transitional changes, for example the decomposition of a residual biomass pool. Inclusion of other data (for example, core-based C accumulation rates from undrained sites or additional flux data from sites that have been rewetted for longer time periods) may help to redress this issue.

Data on net CO₂ fluxes from successfully rewetted tropical organic soils are completely lacking, and data on the CO₂ balance of undrained tropical organic soils are also scarce. Some of the peat swamps in this area may have been degrading naturally for the past 2000 years (Dommain *et al.* 2014). Yet, most tropical peatlands are a net sink of CO₂ (Page *et al.* 2004, Lahteenoja *et al.* 2009, Dommain *et al.* 2011, Lahteenoja *et al.* 2012, Dommain *et al.* 2015), displaying the highest long-term accumulation rates of peatlands worldwide (Dommain *et al.* 2011, Dommain *et al.* 2015). Although the C accumulation data for undrained tropical peatlands (Figure 5) indicate a net sink, the re-establishment of such a sink after rewetting may present a considerable challenge. As such, the default emission factor of 0 t CO₂-C ha⁻¹ yr⁻¹ (Table 2) probably applies to best case rewetting scenarios only.

The removals of CO₂ in rewetted organic soils are in sharp contrast to the high emissions associated with drained organic soils (Table 5 and Figure 6). Moreover, our derived values for DOC losses (Table 3) are lower than those reported for drained organic soils (Evans *et al.* 2015). In addition, N₂O emissions in rewetted soils were very low across all climate zones (Figure 5), although rewetting does result in strongly increased CH₄ emissions following the reversal of drainage (Table 5 and Figure 6b). However, rewetting in general represents a significant climate change mitigation action as a result of (a) the considerable decrease in CO₂ emissions and (b) the accompanying reduction of N₂O emissions (with its high global warming potential) relative to drained organic soils (Table 5 and Figure 6c).

Uncertainties

GHG emissions/removals were characterised by variations both *within* and *between* the disaggregated groups (Figures 4 and 5). Considerable uncertainty is attached to individual data points used in the derivation of the emission factors as most of the studies in the database are generally of a short duration (1–2 years) and do not take into account the longer-term natural variation, a feature captured in long-term datasets (e.g. Roulet *et al.* 2007, McVeigh *et al.* 2014, Aurela *et al.* 2015, Helfter *et al.* 2015). Uncertainty is reduced by using the mean value of multi-year data from the same site and by averaging over multiple sites in each disaggregated group.

The large uncertainties associated with the derived emission factors indicate that individual rewetted and undrained sites may differ considerably in terms of their current abiotic and biotic conditions and resulting vegetation cover. On the one hand, the variation relates to spatial variation found within the various study sites. For example, following rewetting a site may develop as a mosaic of microsites, both vegetated and non-vegetated, characterised by differences in vegetation composition, productivity and *MWTL*, with consequent variations in the magnitude and direction of GHG fluxes (Tuittila *et al.* 1999, Wilson *et al.* 2013). On the other hand, variation between sites can be even larger. Nutrient rich sites in particular display a wider range of flux values between sites than nutrient poor sites (Figure 5), which can be explained by their high hydrological, biogeochemical and botanical diversity. For example, plant associations in rich fens are diverse, ranging from brown moss dominated to sedge stands and reed beds, whereas nutrient poor

temperate and boreal bogs may support a more limited range of plant assemblages. The wide range of flux values in nutrient rich organic soils can also be explained by the diversity of previous land-uses, as nutrient rich organic soils tend to have been used more intensively than nutrient poor sites, especially across the temperate zone (Joosten & Clarke 2002).

The highly generic approach adopted in deriving the IPCC Tier 1 (default) emission factors for rewetted organic soils means that a high level of uncertainty will remain, as is further evidenced by the wide range in the new data values (Figure 5). Clearly, a move towards more specific emission factors (Tier 2) that take into account the factors that control GHG fluxes in rewetted organic soils, and thereby lead to a reduction in the associated uncertainty, is desirable.

Refinement of emission factors

GHG fluxes in rewetted organic soils are controlled by a wide range of external and internal factors, which include the prevailing climate, nutrient status, water table position, previous land use history, time since rewetting, absence or presence of vegetation and vegetation composition. However, in seeking to determine the overarching driver(s) of GHG exchange for rewetted organic soils, the exercise here was constrained to some extent by the quantity of available data for rewetted sites and by the quality of ancillary data in published studies. For the former, we were able to significantly expand the datasets by the inclusion of undrained sites as proxies (Figures 1 and 2), while for the latter it was possible to augment the data in some cases with information contained in other publications from the same site. Our analysis of GHG flux data from rewetted (and undrained) organic soils allowed for an initial disaggregation of the data by climate zone and nutrient status (Figure 3). This is in keeping with the good practices recommended by the IPCC (2006). However, there were insufficient data available to determine the strength of relationships between GHG fluxes and other variables such as previous land use history, time since rewetting and vegetation composition.

Previous land use

Although not captured in the general dataset, the influence of previous land use history (e.g. forest, grassland, cropland and wetland) on GHG fluxes in rewetted organic soils is likely to be profound. For example, CH₄ emissions following the rewetting of former agricultural land can be very high (Hendriks *et al.* 2007, Harpenslager *et al.* 2015) whereas rewetted boreal cutover peatlands may show CH₄

emissions well below the average (Tuittila *et al.* 2000, Waddington & Day 2007). While the influence of previous land use may diminish over time, sub-division of the flux data according to previous land use would, given a sufficient number of studies, undoubtedly improve the accuracy of emissions factors.

Time since rewetting

Available datasets from rewetted organic soils generally cover a period of ten years or less after rewetting and for this reason it is difficult to identify clear temporal patterns in GHG fluxes and determine with accuracy the transition time required to fully capture the changes following rewetting (e.g. Tuittila *et al.* 1999, Bortoluzzi *et al.* 2006, Kivimäki *et al.* 2008, Waddington *et al.* 2010, Wilson *et al.* 2013). Given the limitations in the available scientific literature, our derived emission factors assume that there is no transient period and that rewetted organic soils immediately behave like undrained organic soils in terms of GHG dynamics.

Whereas the high CO₂ emissions observed at drained sites will be reduced immediately upon rewetting, the time needed for recovery of the C sink function may vary from several years to many decades (Tuittila *et al.* 1999, Samaritani *et al.* 2011, Wilson *et al.* 2013) depending on the type of restoration methods employed, how long these methods are continued, and the pre-rewetting climate and hydrological boundary conditions. Re-establishment of the peat-forming vegetation cover on rewetted organic soils is necessary to reinstate the C sink function that ultimately leads to long-term C sequestration in the soil. In the period immediately following rewetting, soil oxidation rates will be low as a consequence of the anoxic conditions, while most of the newly sequestered C is contained within the non-woody biomass pool (leaves, stems, roots and litter). As a result, the ecosystem sink can temporarily be much larger upon restoration (Soini *et al.* 2010, Wilson *et al.* 2013), although a site can remain a CO₂ source during the first years after rewetting as well (Petroni *et al.* 2003, Waddington *et al.* 2010). Over longer time frames (a few decades) a decrease in the amount of CO₂ that is sequestered annually might be expected as the biomass pool eventually approaches a steady state C sequestration rate typical of undrained organic soils.

The impact of rewetting on changes in CH₄ fluxes over time differs between nutrient poor and rich sites. The rewetting of nutrient poor cutover peatlands results in a steady increase in CH₄ emissions in the years immediately after rewetting

as the emerging vegetation cover provides fresh substrates for CH₄ production (Tuittila *et al.* 2000, Waddington & Day 2007). Indeed, work by Vanselow-Algan *et al.* (2015) has indicated that high CH₄ emissions could persist for decades after rewetting, particularly if the vegetation is periodically inundated. In contrast, rewetting of nutrient rich sites seems to result in high initial CH₄ emissions that decline over time as the litter inundated during rewetting activities is decomposed (Limpens *et al.* 2008, Augustin *et al.* 2012). Thus, changes in CH₄ emissions and removals over time appear to be linked to vegetation succession (Tuittila *et al.* 2000, Cooper *et al.* 2014) and understanding how emissions change over time would require the inclusion of vegetation information.

Vegetation composition

Numerous studies have indicated the important role of vegetation in the regulation of GHG fluxes in both undrained and rewetted organic soils. Plants remove CO₂ from the atmosphere *via* photosynthesis, utilise available nitrogen in the soil, and provide the substrate for CH₄ production and a pathway for the transportation of CH₄ from the saturated soil to the atmosphere (e.g. Bubier 1995, Shannon *et al.* 1996, Tuittila *et al.* 2000, Marinier *et al.* 2004, Wilson *et al.* 2009, Dias *et al.* 2010). Indeed, the presence of shunt species (i.e. wetland adapted vascular plant species known to transport CH₄ from the soil to the atmosphere) has a significant effect on CH₄ efflux from organic soils (e.g. Couwenberg & Fritz 2012, Levy *et al.* 2012) and a refinement of emission factors could be achieved through the development of nationally or regionally specific emission factors that directly address vegetation composition (see Riutta *et al.* 2007, Dias *et al.* 2010, Couwenberg *et al.* 2011, Forbrich *et al.* 2011). In particular, where perennial woody biomass plays a significant role in the net CO₂ and CH₄ exchange (e.g. pneumatophore species in the tropics) between rewetted organic soils and the atmosphere (e.g. Pangala *et al.* 2013), country-specific methods should be developed that reflect the C stock changes in the tree biomass and dead tree organic matter pools under typical management practices and their interaction with the soil pool.

Water table level

The relationship between water table level and CO₂/CH₄ emissions/removals was evident in this study (Figures 1 and 2). As the water table is one of the main controls on GHG cycling, future (i.e. country specific) emission factors could be derived

and disaggregated by water table level provided sufficient ancillary data are available (e.g. mean annual, maximum and minimum water table values). Drainage ditches have been shown to be “hotspots” of CH₄ emissions within the wider drained landscape (Cooper *et al.* 2014, IPCC 2014a). Few data are available on CH₄ emissions from ditches that remain after rewetting or that are filled in during rewetting activities, although there is some evidence to suggest that CH₄ emissions may remain high after rewetting (Waddington & Day 2007, Cooper *et al.* 2014). However, rewetting reduces the hydrological differences between fields and neighbouring ditches creating a more homogeneous surface that is not so different from undrained sites where hollows are major hotspots of CH₄ emissions (e.g. A. Laine *et al.* 2007b). Improved estimates of water table level distribution across a site would better capture the spatial variability associated with CH₄ fluxes. Our literature search also identified the impact of inundation on GHG dynamics in rewetted sites. In many cases, where the water table is maintained at very high levels (>20cm above the soil surface), CH₄ emissions can be extremely high (e.g. Augustin & Chojnicki 2008, Koch *et al.* 2014, Hahn *et al.* 2015, Vanselow-Algan *et al.* 2015), although much lower values have been observed as well (Koch *et al.* 2014, Minke *et al.* 2015). More research will be needed to assess the drivers behind the wide variation found in essentially flooded ecosystems.

Improved data collection

Emission factors could be further refined by the use of advanced process-based (mechanistic) models. Annual GHG fluxes are commonly calculated using quasi-mechanistic models that rely on descriptive attributes (*WTD*, temperature, photon flux density, vegetation cover, *etc.*) fitted to intermittent measurement data (Minke *et al.* 2012). Process-based models have the potential to integrate the interactions between biomass, dead organic matter and soil carbon pools, and to provide improved spatial and temporal estimates of GHG exchange; they do require, however, a very high level of information and complexity in regard to the interactions and processes described above (e.g. Walter *et al.* 2001, Frohling *et al.* 2002, Li *et al.* 2010, Baird *et al.* 2012, Meng *et al.* 2012, Gong *et al.* 2013, Metzger *et al.* 2015). Furthermore, the use of more sophisticated models would not remove the need for robust field measurements, which are required to support model development, parameterisation and testing.

More refined emission factors of course need to

be accompanied by equally disaggregated land cover data. Recent advances in high resolution remote sensing and aerial imagery have the potential to significantly improve data collection. For example, unmanned aerial vehicles (UAVs) have been shown to provide extremely high resolution imagery in terms of the areal cover of vegetation communities (Kalacska *et al.* 2013) and could be particularly useful for sites where there is a mosaic of microsites (e.g. Knoth *et al.* 2013). Moreover, the availability of new satellite platforms, such as the Sentinel 2B (<https://earth.esa.int/web/guest/missions/esa-operational-eo-missions/sentinel-2>), could provide highly detailed imagery that would allow further disaggregation by vegetation type at a relatively low cost. Similarly, aerial platforms could potentially provide more detailed information to allow disaggregation by soil moisture or water table level (Kasischke *et al.* 2009, Jaenicke *et al.* 2011, Torbick *et al.* 2012).

Information gaps and ongoing issues

While the number of GHG studies on organic soils has steadily increased over the last few decades (see reviews by Lai 2009, Haddaway *et al.* 2014, IPCC 2014a), our work here has highlighted the existence of some research gaps, in particular the paucity of GHG data from the tropical climate zones as well as specific issues associated with rewetting.

Tropical data

GHG emissions from drained tropical peatlands are affected by a number of factors, such as higher peat surface temperatures following removal of the forest vegetation, aerobic conditions in the upper part of the peat column resulting from both drainage and disturbance caused by management operations and fertiliser applications (Jauhiainen *et al.* 2012, Jauhiainen *et al.* 2014). The effect of rewetting on GHG fluxes from tropical organic soils has yet to be measured. If the soil becomes persistently waterlogged, CO₂ fluxes should be near zero from a biochemical perspective. However, it is much more difficult to rewet and maintain a stable water table in tropical peat than in boreal and temperate peat soils because the hydraulic conductivity of tropical peat is extremely high (Page *et al.* 2008). Furthermore, there is a tendency to flooding during the wet season and near drought in severely dry periods such as El Niño years (Dommain *et al.* 2011). Moreover, several studies in south-east Asia have indicated that there could be sizeable C emissions from former agricultural lands even if these are essentially wet (Hooijer *et al.* 2012, Jauhiainen *et al.* 2012, Husnain *et al.* 2014), while Gandois *et al.* (2013) reported

substantial impacts on the C balance of an undrained tropical peat swamp forest following deforestation alone. The inference from these studies is that it may take considerable time for C losses from tropical peat swamp forests to reduce after rewetting. Moreover, achieving pre-disturbance C accumulation rates will be a long and largely unpredictable process that will probably only be realised following the re-establishment of closed-canopy forest, along with the accompanying environmental conditions conducive to peat formation. Additional research on rewetting techniques and associated GHG fluxes will be needed as a basis for higher tier emission factors. Furthermore, reliable data on GHG fluxes from organic soils in Africa, the tropical Americas or other tropical regions outside south-east Asia are very rare and virtually nothing is known about the effects of rewetting.

Fire

Due to high moisture contents, organic soils in intact ecosystems are protected to some degree from burning (Turetsky *et al.* 2015), although fires do occur on undrained organic soils. It can be assumed that the probability of fire occurrence in rewetted organic soils is likely to be small if the water table position is maintained at or near the surface, although there may still be a major risk of fire affecting trees in tropical regions where the temperature is high, even if the water table is near the peat surface. If the surface peat does become dry and flammable, for example during periods of drought, wet layers deeper in the peat profile will serve as fire barriers, limiting the depth of peat burning and hence C loss.

Waterborne pathways

An understanding of the fate of DOC leaked from organic soils is still poor. While DOC can be returned to the atmosphere as CO₂ (or CH₄), or transferred to lake sediments or long term C stores such as the deep ocean or marine sediments (Müller *et al.* 2015, Abrams *et al.* 2016), further studies should be carried out to improve the values for the conversion factor $Frac_{DOC_{CO_2}}$. Measurements from undrained and rewetted organic soils should be undertaken to obtain more accurate and country-specific values of DOC_{FLUX}. Furthermore, since DOC production has been observed to vary with vegetation composition and productivity as well as with soil temperature, research should focus on developing specific values for different types of rewetted organic soils (e.g. nutrient rich *versus* nutrient poor). Additional waterborne C losses come

from leaching of inorganic carbon (DIC) and from erosion (particulate organic carbon; POC), as well as evasion of CO₂ and CH₄ from the waterbodies within or close to the peatlands themselves. However, these are not treated in the *Wetlands Supplement* as very few data exist from rewetted sites and these losses are likely to be site specific (Evans *et al.* 2015). Although, compared to DOC, a greater proportion of POC may be simply translocated from the rewetted organic soil to other stable C stores such as freshwater or marine sediments (where it will not lead to CO₂ emissions), there is now some evidence to suggest that POC may also undergo significant mineralisation during riverine transport or (since POC fluxes are typically highest during flood events) following overbank deposition onto floodplains (Evans *et al.* 2015).

CONCLUSIONS

Rewetting of organic soils results in a decrease in CO₂ and N₂O emissions, DOC losses and GHG emissions based on the global warming potential; but also leads to an increase in CH₄ emissions. In general, the emission factors derived in the *Wetlands Supplement* for rewetted organic soils appear robust and compare well to the new data (n=58) published since the Supplement. However, the GHG emission factor estimates derived in this paper are subject to uncertainty as a consequence of the lack of data for some climate zones and land use categories, and the wide variation in GHG emissions/removals inherent to organic soils in general. Future research should focus on the information gaps that have been highlighted here, with particular emphasis on determining the transient period following rewetting, reducing uncertainty through the derivation of country specific emission factors and, perhaps most critically given the high emissions associated with drainage, the quantification of GHG emissions/removals in rewetted tropical organic soils.

ACKNOWLEDGEMENTS

We would like to acknowledge the inputs of our colleagues Maya Fukuda, Osamu Nagata and Faizal Parish to the derivation of the *Wetlands Supplement* emission factors. We would also like to thank the Editor and two anonymous reviewers for their insightful comments and suggestions on earlier drafts of this article.

REFERENCES

- Abrams, J.F., Hohn, S., Rixen, T., Baum, A. & Merico, A. (2016) The impact of Indonesian peatland degradation on downstream marine ecosystems and the global carbon cycle. *Global Change Biology*, 22(1), 325–337.
- Adji, F.F., Hamada, Y., Darang, U., Limin, S.H. & Hatano, R. (2014) Effect of plant-mediated oxygen supply and drainage on greenhouse gas emission from a tropical peatland in Central Kalimantan, Indonesia. *Soil Science and Plant Nutrition*, 60(2), 216–230.
- Adkinson, A.C., Syed, K.H. & Flanagan, L.B. (2011) Contrasting responses of growing season ecosystem CO₂ exchange to variation in temperature and water table depth in two peatlands in northern Alberta, Canada. *Journal of Geophysical Research*, 116(G1), G01004.
- Ågren, A., Jansson, M., Ivarsson, H., Bishop, K. & Seibert, J. (2008) Seasonal and runoff-related changes in total organic carbon concentrations in the River Öre, Northern Sweden. *Aquatic Sciences*, 70(1), 21–29.
- Alkhatib, M., Jennerjahn, T.C. & Samiaji, J. (2007) Biogeochemistry of the Dumai River estuary, Sumatra, Indonesia, a tropical black-water river. *Limnology and Oceanography*, 52(6), 2410–2417.
- Alm, J., Shurpali, N.J., Tuittila, E.-S., Laurila, T., Maljanen, M., Saarnio, S. & Minkkinen, K. (2007) Methods for determining emission factors for the use of peat and peatlands - flux measurements and modelling. *Boreal Environment Research*, 12, 85–100.
- Alm, J., Talanov, A., Saarnio, S., Silvola, J., Ikkonen, E., Aaltonen, H., Nykänen, H. & Martikainen, P.J. (1997) Reconstruction of the carbon balance for microsites in a boreal oligotrophic pine fen, Finland. *Oecologia*, 110, 423–431.
- Augustin, J. (2003) Gaseous emissions from constructed wetlands and (re)flooded meadows. *Publicationes Instituti Geographici Universitatis Tartuensis*, 94, 3–8.
- Augustin, J. & Chojnicki, B. (2008) Austausch von klimarelevanten Spurengasen, Klimawirkung und Kohlenstoffdynamik in den ersten Jahren nach der Wiedervernässung von degradiertem Niedermoorgrünland (Exchange of climate-relevant trace gases, climate effect and carbon dynamics in the first years after rewetting of degraded fen grassland). In: Gelbrecht J., Zak D. & Augustin J. (eds.) *Phosphor- und Kohlenstoff-Dynamik und Vegetationsentwicklung in wiedervernässten Mooren des Peenetales in Mecklenburg-Vorpommern - Status, Steuergrößen und Handlungsmöglichkeiten (Phosphorus and Carbon Dynamics and Vegetation Development in Rewetted Fens of the Peene Valley in Mecklenburg-Western Pomerania - Status, Control Variables and Possibilities for Action)*, Berichte des IGB 26, Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, 50–67 (in German).
- Augustin, J., Couwenberg, J. & Minke, M. (2012) Peatlands and greenhouse gases. In: Tanneberger, F. & Wichtmann, W. (eds.) *Carbon Credits From Peatlands Rewetting. Climate - Biodiversity - Land Use*, Schweizerbart, Stuttgart, 13–19.
- Augustin, J. & Merbach, W. (1998) Greenhouse gas emissions from fen mires in northern Germany: quantification and regulation. In: Merbach, W. & Wittenmayer, L. (eds.) *Beiträge aus der Hallenser Pflanzenernährungsforschung (Contributions from the Halle Plant Nutrition Research)*, Grauer, Beuren, 97–110.
- Augustin, J., Merbach, W., Käding, H., Schmidt, W. & Schalitz, G. (2006) Lachgas- und Methanemission aus degradierten Niedermoorstandorten Nordostdeutschlands unter dem Einfluß unterschiedlicher Bewirtschaftung (Nitrous oxide and methane emissions from degraded fens in north-east Germany under the influence of different management). In: Alfred-Wegener-Stiftung (ed.) *Von den Ressourcen zum Recycling (From Resources To Recycling)*, Verlag Ernst & Sohn, Berlin, 131–139 (in German).
- Aurela, M., Laurila, T. & Tuovinen, J.-P. (2002) Annual CO₂ balance of a subarctic fen in northern Europe: Importance of the wintertime efflux. *Journal of Geophysical Research*, 107(D21), 4607.
- Aurela, M., Lohila, A., Tuovinen, J.-P., Hatakka, J., Penttilä, T. & Laurila, T. (2015) Carbon dioxide and energy flux measurements in four northern-boreal ecosystems at Pallas. *Boreal Environment Research*, 20, 455–473.
- Aurela, M., Lohila, A., Tuovinen, J.-P., Hatakka, J., Riutta, T. & Laurila, T. (2009) Carbon dioxide exchange on a northern boreal fen. *Boreal Environment Research*, 14, 699–710.
- Baird, A., Morris, P. & Belyea, L.R. (2012) The DigiBog peatland development model 1: rationale, conceptual model and hydrological basis. *Ecohydrology*, 5, 242–255.
- Ballantyne, D., Hrilbljan, J., Pypker, T. & Chimner,

- R.A. (2014) Long-term water table manipulation alters peatlands gaseous carbon fluxes in Northern Michigan. *Wetlands Ecology and Management*, 22, 35–47.
- Bartlett, K.B. & Harriss, R.C. (1993) Review and assessment of methane emissions from wetlands. *Chemosphere*, 26(1–4), 261–320.
- Baum, A., Rixen, T. & Samiaji, J. (2007) Relevance of peat draining rivers in central Sumatra for the riverine input of dissolved organic carbon into the ocean. *Estuarine, Coastal and Shelf Science*, 73(3–4), 563–570.
- Beetz, S., Liebersbach, H., Glatzel, S., Jurasinski, G., Buczko, U. & Höper, H. (2013) Effects of land use intensity on the full greenhouse gas balance in an Atlantic peat bog. *Biogeosciences*, 10, 1067–1082.
- Beyer, C. & Höper, H. (2015) Greenhouse gas exchange of rewetted bog peat extraction sites and a *Sphagnum* cultivation site in northwest Germany. *Biogeosciences*, 12(7), 2101–2117.
- Billett, M.F., Charman, D.J., Clark, J.M., Evans, C.D., Evans, M.G., Ostle, N.J., Worrall, F., Burden, A., Dinsmore, K.J., Jones, T., McNamara, N.P., Parry, L., Rowson, J.G. & Rose, R. (2010) Carbon balance of UK peatlands: current state of knowledge and future research challenges. *Climate Research*, 45, 13–29.
- Billett, M.F., Palmer, S.M., Hope, D., Deacon, C., Storeton-West, R., Hargreaves, K.J., Flechard, C. & Fowler, D. (2004) Linking land-atmosphere-stream carbon fluxes in a lowland peatland system. *Global Biogeochemical Cycles*, 18(1), GB1024.
- Bortoluzzi, E., Epron, D., Siegenthaler, A., Gilbert, D. & Buttler, A. (2006) Carbon balance of a European mountain bog at contrasting stages of regeneration. *New Phytologist*, 172(4), 708–718.
- Bubier, J.L. (1995) The relationship of vegetation to methane emission and hydrochemical gradients in northern peatlands. *Journal of Ecology*, 83, 403–420.
- Bubier, J.L., Frolking, S., Crill, P.M. & Linder, E. (1999) Net ecosystem productivity and its uncertainty in a diverse boreal peatland. *Journal of Geophysical Research*, 104(D22), 27683–27692.
- Bubier, J.L., Moore, T.R. & Roulet, N.T. (1993) Methane emissions from wetlands in the mid-boreal region of northern Ontario, Canada. *Ecology*, 74(8), 2240–2254.
- Cagampan, J. & Waddington, J.M. (2008) Net ecosystem CO₂ exchange of a cutover peatland rehabilitated with a transplanted acrotelm. *Ecoscience*, 15(2), 258–267.
- Campbell, D.I., Smith, J., Goodrich, J.P., Wall, A.M. & Schipper, L.A. (2014) Year-round growing conditions explains large CO₂ sink strength in a New Zealand raised peat bog. *Agricultural and Forest Meteorology*, 192–193, 59–68.
- Chimner, R.A. & Karberg, J.M. (2008) Long-term carbon accumulation in two tropical mountain peatlands, Andes Mountains, Ecuador. *Mires and Peat*, 3(04), 1–10.
- Chistotin, M.V., Sirin, A.A. & Dulov, L.E. (2006) Sezonnaja dinamika jemissii uglekislogo gaza i metana pri osushenii bolota v Moskovskoj oblasti dlja dobychi torfa i sel'skohozjajstvennogo ispol'zovanija (Seasonal dynamics of carbon dioxide and methane emission from a peatland in Moscow Region drained for peat extraction and agricultural use). *Agrohimiya (Agrochemistry)*, 6, 54–62 (in Russian). Abstract online at: <http://elibrary.ru/item.asp?id=9241689>, accessed 31 Mar 2016.
- Christensen, T.R., Jackowicz-Korczyński, M., Aurela, M., Crill, P., Heliasz, M., Mastepanov, M. & Friborg, T. (2012) Monitoring the multi-year carbon balance of a subarctic palsa mire with micrometeorological techniques. *Ambio*, 41(3), 207–217.
- Clair, T.A., Arp, P., Moore, T.R., Dalvac, M. & Meng, F.-R. (2002) Gaseous carbon dioxide and methane, as well as dissolved organic carbon losses from a small temperate wetland under a changing climate. *Environmental Pollution*, 116, Suppl 1, S143–S148.
- Cleary, J., Roulet, N.T. & Moore, T.R. (2005) Greenhouse gas emissions from Canadian peat extraction, 1990–2000: a life cycle analysis. *Ambio*, 34(6), 456–461.
- Cliche Trudeau, N., Garneau, M. & Pelletier, L. (2013) Methane fluxes from a patterned fen of the northeastern part of the La Grande river watershed, James Bay, Canada. *Biogeochemistry*, 113(1–3), 409–422.
- Cliche Trudeau, N., Garneau, M. & Pelletier, L. (2014) Interannual variability in the CO₂ balance of a boreal patterned fen, James Bay, Canada. *Biogeochemistry*, 118(1–3), 371–387.
- Clymo, R.S. & Reddaway, E.J.F. (1971) Productivity of *Sphagnum* (bog-moss) and peat accumulation. *Hidrobiologia* (Bucharest), 12, 181–192. (See: Clymo, R.S. & Reddaway, E.J.F. (1972) A tentative dry-matter balance sheet for the wet blanket bog on Burnt Hill, Moor House NNR. *Aspects of the Ecology of the Northern Pennines, Occasional Papers*, 3, Institute of

- Terrestrial Ecology, Moor House, 1–15).
- Cooper, M.D.A., Evans, C.D., Zielinski, P., Levy, P.E., Gray, A., Peacock, M., Norris, D., Fenner, N. & Freeman, C. (2014) Infilled ditches are hotspots of landscape methane flux following peatland re-wetting. *Ecosystems*, 17(7), 1227–1241.
- Couwenberg, J., Dommain, R. & Joosten, H. (2010) Greenhouse gas fluxes from tropical peatlands in south-east Asia. *Global Change Biology*, 16(6), 1715–1732.
- Couwenberg, J. & Fritz, C. (2012) Towards developing IPCC methane ‘emission factors’ for peatlands (organic soils). *Mires and Peat*, 10(03), 1–17.
- Couwenberg, J., Thiele, A., Tanneberger, F., Augustin, J., Bärtsch, S., Dubovik, D., Liashchynskaya, N., Michaelis, D., Minke, M., Skuratovich, A. & Joosten, H. (2011) Assessing greenhouse gas emissions from peatlands using vegetation as a proxy. *Hydrobiologia*, 674(1), 67–89.
- Dawson, J.J.C., Billet, M., Hope, D., Palmer, S.M. & Deacon, C. (2004) Sources and sinks of aquatic carbon in a peatland stream continuum. *Biogeochemistry*, 70, 71–92.
- Dias, A., Hoorens, B., Van Logtestijn, R., Vermaat, J. & Aerts, R. (2010) Plant species composition can be used as a proxy to predict methane emissions in peatland ecosystems after land-use changes. *Ecosystems*, 13(4), 526–538.
- di Folco, M.-B. & Kirkpatrick, J.B. (2011) Topographic variation in burning-induced loss of carbon from organic soils in Tasmanian moorlands. *Catena*, 87(2), 216–225.
- Dinsmore, K.J., Smart, R.P., Billett, M.F., Holden, J., Baird, A.J. & Chapman, P.J. (2011) Greenhouse gas losses from peatland pipes: a major pathway for loss to the atmosphere? *Journal of Geophysical Research*, 116(G3), doi:10.1029/2011JG001646.
- Dise, N.B., Gorham, E. & Verry, E.S. (1993) Environmental factors controlling methane emissions from peatlands in northern Minnesota. *Journal of Geophysical Research: Atmospheres*, 98(D6), 10583–10594.
- Dommain, R., Cobb, A.R., Joosten, H., Glaser, P.H., Chua, A.F.L., Gandois, L., Kai, F.-M., Noren, A., Salim, K.A., Su’ut, N.S.H. & Harvey, C.F. (2015) Forest dynamics and tip-up pools drive pulses of high carbon accumulation rates in a tropical peat dome in Borneo (Southeast Asia). *Journal of Geophysical Research: Biogeosciences*, 120(4), 617–640.
- Dommain, R., Couwenberg, J., Glaser, P.H., Joosten, H. & Suryadiputra, I.N.N. (2014) Carbon storage and release in Indonesian peatlands since the last deglaciation. *Quaternary Science Reviews*, 97, 1–32.
- Dommain, R., Couwenberg, J. & Joosten, H. (2011) Development and carbon sequestration of tropical peat domes in south-east Asia: links to post-glacial sea-level changes and Holocene climate variability. *Quaternary Science Reviews*, 30(7–8), 999–1010.
- Drewer, J., Lohila, A., Aurela, M., Laurila, T., Minkinen, K., Penttilä, T., Dinsmore, K.J., McKenzie, R.M., Helfter, C., Flechard, C., Sutton, M.A. & Skiba, U.M. (2010) Comparison of greenhouse gas fluxes and nitrogen budgets from an ombrotrophic bog in Scotland and a minerotrophic sedge fen in Finland. *European Journal of Soil Science*, 65(5), 640–650, doi:10.1111/j.1365-2389.2010.01267.x.
- Drösler, M. (2005) Trace gas exchange and climatic relevance of bog ecosystems, southern Germany. PhD thesis, Universität München, Germany, 182 pp.
- Drösler, M., Freibauer, A., Adelman, W., Augustin, J., Bergman, L., Beyer, C., Chojnicki, B.H., Förster, C., Giebels, M., Görlitz, S., Höper, H., Kantelhardt, J., Liebersbach, H., Hahn-Schöfl, M., Minke, M., Petschow, U., Pfadenhauer, J., Schaller, L., Schägner, P., Sommer, M., Thuile, A. & Wehrhan, M. (2013) *Klimaschutz durch Moorschutz in der Praxis (Climate Protection through Peatland Protection in Practice)*. Institut für Agrarrelevante Klimaforschung (AK), Johann Heinrich von Thünen-Institut, Braunschweig, 15 pp. (in German with English summary).
- Ernfors, M., von Arnold, K., Stendahl, J., Olsson, M. & Klemetsson, L. (2008) Nitrous oxide emissions from drained organic forest soils—an up-scaling based on C:N ratios. *Biogeochemistry*, 89, 29–41.
- Evans, C., Renou-Wilson, F. & Strack, M. (2015) The role of waterborne carbon in the greenhouse gas balance of drained and re-wetted peatlands. *Aquatic Sciences*, doi:10.1007/s00027-015-0447-y, 18 pp.
- Fiedler, S. & Sommer, M. (2000) Methane emissions, groundwater levels and redox potentials of common wetland soils in a temperate-humid climate. *Global Biogeochemical Cycles*, 14(4), 1081–1093.
- Flessa, H., Wild, U., Klemisch, M. & Pfadenhauer, J. (1997) C- und N-Stoffflüsse auf Torfstichsimulationsflächen im Donaumoos (C and N matter flow on simulated peat cutting

- areas in the Donaumoos). *Zeitschrift für Kulturtechnik und Landentwicklung (Journal for Cultivation Techniques and Land Development)*, 38, 11–17 (in German).
- Forbrich, I., Kutzbach, L., Wille, C., Becker, T., Wu, J. & Wilmking, M. (2011) Cross-evaluation of measurements of peatland methane emissions on microform and ecosystem scales using high-resolution landcover classification and source weight modelling. *Agricultural and Forest Meteorology*, 151(7), 864–874.
- Frolking, S., Roulet, N.T., Moore, T.R., Lafleur, P.M., Bubier, J.L. & Crill, P.M. (2002) Modeling seasonal to annual carbon balance of Mer Bleue Bog, Ontario, Canada. *Global Biogeochemical Cycles*, 16(3), 4-1–4-21.
- Furukawa, Y., Inubushi, K., Ali, M., Itang, A.M. & Tsuruta, H. (2005) Effect of changing groundwater levels caused by land-use changes on greenhouse gas fluxes from tropical peat lands. *Nutrient Cycling in Agroecosystems*, 71(1), 81–91.
- Gandois, L., Cobb, A.R., Hei, I.C., Lim, L.B.L., Salim, K.A. & Harvey, C.F. (2013) Impact of deforestation on solid and dissolved organic matter characteristics of tropical peat forests: implications for carbon release. *Biogeochemistry*, 114(1–3), 183–199.
- Gauci, V., Dise, N.B. & Fowler, D. (2002) Controls on suppression of methane flux from a peat bog subjected to simulated acid rain sulfate deposition. *Global Biogeochemical Cycles*, 16(1), 4-1–4-12, doi:10.1029/2000GB001370.
- Gažovič, M., Forbrich, I., Jager, D.F., Kutzbach, L., Wille, C. & Wilmking, M. (2013) Hydrology-driven ecosystem respiration determines the carbon balance of a boreal peatland. *Science of the Total Environment*, 463–464, 675–682.
- Glatzel, S., Koebsch, F., Beetz, S., Hahn, J., Richter, P. & Jurasinski, G. (2011) Maßnahmen zur Minderung der Treibhausgas Freisetzung aus Mooren im Mittleren Mecklenburg (Measures to lower greenhouse gas emissions from peatlands in central Mecklenburg). *Telma*, 4, 85–106 (in German).
- Golovatskaya, E. & Dyukarev, E. (2009) Carbon budget of oligotrophic mire sites in the Southern Taiga of Western Siberia. *Plant and Soil*, 315(1), 19–34.
- Gong, J., Kellomäki, S., Wang, K., Zhang, C., Shurpali, N. & Martikainen, P.J. (2013) Modeling CO₂ and CH₄ flux changes in pristine peatlands of Finland under changing climate conditions. *Ecological Modelling*, 263, 64–80.
- Goodrich, J.P., Campbell, D.I., Roulet, N.T., Clearwater, M.J. & Schipper, L.A. (2015) Overriding control of methane flux temporal variability by water table dynamics in a Southern Hemisphere, raised bog. *Journal of Geophysical Research: Biogeosciences*, 120 (5), 819–831
- Günther, A.B., Huth, V., Jurasinski, G. & Glatzel, S. (2014) Scale-dependent temporal variation in determining the methane balance of a temperate fen. *Greenhouse Gas Measurement and Management*, 4(1), 41–48.
- Günther, A., Huth, V., Jurasinski, G. & Glatzel, S. (2015) The effect of biomass harvesting on greenhouse gas emissions from a rewetted temperate fen. *Global Change Biology: Bioenergy*, 7(4), 1092–1106.
- Haddaway, N.R., Burden, A., Evans, C.D., Healey, J.R., Jones, D.L., Dalrymple, S.E. & Pullin, A.S. (2014) Evaluating effects of land management on greenhouse gas fluxes and carbon balances in boreo-temperate lowland peatland systems. *Environmental Evidence*, 3(5), doi:10.1186/2047-2382-3-5.
- Hadi, A., Haradi, M., Inubushi, K., Purnomo, E., Razie, F. & Tsuruta, H. (2001) Effects of land-use change in tropical peat soil on the microbial population and emission of greenhouse gases. *Microbes and Environment*, 16(2), 79–86.
- Hadi, A., Inubushi, K., Furukawa, Y., Purnomo, E., Rasmadi, M. & Tsuruta, H. (2005) Greenhouse gas emissions from tropical peatlands of Kalimantan, Indonesia. *Nutrient Cycling in Agroecosystems*, 71(1), 73–80.
- Hahn, J., Köhler, S., Glatzel, S. & Jurasinski, G. (2015) Methane exchange in a coastal fen in the first year after flooding - a systems shift. *PLoS ONE*, 10(10), e0140657.
- Harazono, Y., Mano, M., Miyata, A., Zulueta, R. & Oechel, W.C. (2003) Inter-annual carbon dioxide uptake of a wet sedge tundra ecosystem in the Arctic. *Tellus*, 55B, 215–231.
- Harpenslager, S.F., van den Elzen, E., Kox, M.A.R., Smolders, A.J.P., Ettwig, K.F. & Lamers, L.P.M. (2015) Rewetting former agricultural peatlands: Topsoil removal as a prerequisite to avoid strong nutrient and greenhouse gas emissions. *Ecological Engineering*, 84, 159–168.
- Heikkinen, J.E.P., Elsakov, V. & Martikainen, P.J. (2002) Carbon dioxide and methane dynamics and annual carbon balance in tundra wetland in NE Europe, Russia. *Global Biogeochemical Cycles*, 16(4), 1115.
- Helfter, C., Campbell, C., Dinsmore, K.J., Drewer, J., Coyle, M., Anderson, M., Skiba, U., Nemitz, E., Billett, M.F. & Sutton, M.A. (2015) Drivers of long-term variability in CO₂ net ecosystem

- exchange in a temperate peatland. *Biogeosciences*, 12(6), 1799–1811.
- Hendriks, D.M.D., van Huissteden, J., Dolman, A.J. & van den Molen, M.K. (2007) The full greenhouse gas balance of an abandoned peat meadow. *Biogeosciences*, 4, 411–424.
- Herbst, M., Friborg, T., Schelde, K., Jensen, R., Ringgaard, R., Vasquez, V., Thomsen, A.G. & Soegaard, H. (2013) Climate and site management as driving factors for the atmospheric greenhouse gas exchange of a restored wetland. *Biogeosciences*, 10, 39–52.
- Hirano, T., Jauhiainen, J., Inoue, T. & Takahashi, H. (2009) Controls on the carbon balance of tropical peatlands. *Ecosystems*, 12(6), 873–887.
- Hooijer, A., Page, S., Jauhiainen, J., Lee, W.A., Lu, X.X., Idris, A. & Anshari, C. (2012) Subsidence and carbon loss in drained tropical peatlands. *Biogeosciences*, 9, 1053–1071.
- Hribljan, J.A., Cooper, D.J., Sueltenfuss, J., Wolf, E.C., Heckman, K.A., Lilleskov, E.A. & Chimner, R.A. (2015) Carbon storage and long-term rate of accumulation in high-altitude Andean peatlands of Bolivia. *Mires and Peat*, 15(12), 1–14.
- Humphreys, E.R., Charron, C., Brown, M. & Jones, R. (2014) Two bogs in the Canadian Hudson Bay Lowlands and a temperate bog reveal similar annual net ecosystem exchange of CO₂. *Arctic, Antarctic, and Alpine Research*, 46(1), 103–113.
- Husnain, H., Wigena, I.G.P., Dariah, A., Marwanto, S., Setyanto, P. & Agus, F. (2014) CO₂ emissions from tropical drained peat in Sumatra, Indonesia. *Mitigation and Adaptation Strategies for Global Change*, 19(6), 845–862.
- Huth, V., Günther, A., Jurasinski, G. & Glatzel, S. (2013) The effect of an exceptionally wet summer on methane effluxes from a 15-year rewetted fen in north-east Germany. *Mires and Peat*, 13(02), 1–7.
- Inubushi, K., Furukawa, Y., Hadi, A., Purnomo, E. & Tsuruta, H. (2003) Seasonal changes of CO₂, CH₄ and N₂O fluxes in relation to land-use change in tropical peatlands located in coastal area of South Kalimantan. *Chemosphere*, 52(3), 603–608.
- Inubushi, K., Hadi, A., Okazaki, M. & Yonebayashi, K. (1998) Effect of converting wetland forest to sage palm plantations on methane gas flux and organic carbon dynamics in tropical peat soil. *Hydrological Processes*, 12(13–14), 2073–2080.
- IPCC (1996) *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories* (3 volumes). Online at: <http://www.ipcc-nggip.iges.or.jp/public/gl/invs1.html>, accessed 23 Mar 2016.
- IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (5 volumes). Prepared by the National Greenhouse Gas Inventories Programme. Eggleston, H.S., Buendia, L., Miwa, K., Ngara, T. & Tanabe, K. (eds.), IGES, Japan. Online at: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html>, accessed 23 Mar 2016.
- IPCC (2014a) *2013 Supplement to the 2006 Intergovernmental Panel on Climate Change Guidelines for National Greenhouse Gas Inventories: Wetlands*. Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. & Troxler, T.G. (eds.), IPCC, Switzerland. Online at: <http://www.ipcc-nggip.iges.or.jp/public/wetlands/index.html>, accessed 23 Mar 2016.
- IPCC (2014b) *2013 Revised Supplementary Methods and Good Practice Guidance Arising from the Kyoto Protocol*. Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. & Troxler, T.G. (eds.), IPCC, Switzerland. Online at: <http://www.ipcc-nggip.iges.or.jp/public/kpsg/index.html>, accessed 23 Mar 2016.
- Jacobs, C.M.J., Jacobs, A.F.G., Bosveld, F.C., Hendriks, D.M.D., Hensen, A., Kroon, P.S., Moors, E., Nol, I., Schrier-Uijl, A. & Veenendaal, E.M. (2007) Variability of annual CO₂ exchange from Dutch grasslands. *Biogeosciences*, 4, 803–816.
- Jaenicke, J., Enghart, S. & Siegert, F. (2011) Monitoring the effect of restoration measures in Indonesian peatlands by radar satellite imagery. *Journal of Environmental Management*, 92(3), 630–638.
- Jager, D.F., Wilmking, M. & Kukkonen, J. (2009) The influence of summer seasonal extremes on dissolved organic carbon export from a boreal peatland catchment: evidence from one dry and one wet growing season. *Science of the Total Environment*, 407, 1373–1382.
- Jauhiainen, J., Heikkinen, J., Martikainen, P.J. & Vasander, H. (2001) CO₂ and CH₄ fluxes in pristine peat swamp forest and peatland converted to agriculture in Central Kalimantan, Indonesia. *International Peat Journal*, 11, 43–49.
- Jauhiainen, J., Hooijer, A. & Page, S.E. (2012) Carbon dioxide emissions from an *Acacia* plantation on peatland in Sumatra, Indonesia. *Biogeosciences*, 9(2), 617–630.
- Jauhiainen, J., Kerojoki, O., Silvennoinen, H., Limin, S. & Vasander, H. (2014) Heterotrophic respiration in drained tropical peat is greatly

- affected by temperature—a passive ecosystem cooling experiment. *Environmental Research Letters*, 9, 105013, 18 pp. doi:10.1088/1748-9326/9/10/105013.
- Jauhiainen, J., Limin, S., Silvennoinen, H. & Vasander, H. (2008) Carbon dioxide and methane fluxes in drained tropical peat before and after hydrological restoration. *Ecology*, 89(12), 3503–3514.
- Jauhiainen, J. & Silvennoinen, H. (2012) Diffusion GHG fluxes at tropical peatland drainage canal water surfaces. *Suo*, 63(3–4), 93–105.
- Jauhiainen, J., Takahashi, H., Heikkinen, J.E.P., Martikainen, P.J. & Vasander, H. (2005) Carbon fluxes from a tropical peat swamp forest floor. *Global Change Biology* 11(10), 1788–1797.
- Jauhiainen, J., Vasander, H., Jaya, A., Inoue, T., Heikkinen, J. & Martikainen, P. (2004) Carbon balance in managed tropical peat in Central Kalimantan, Indonesia. In: Päivänen J. (ed.) *Wise use of Peatlands, Proceedings of the 12th International Peat Congress*, International Peat Society, Tampere, Finland, 653–659.
- Joosten, H. & Clarke, D. (2002) *Wise Use of Mires and Peatlands: Background and Principles Including a Framework for Decision-making*. International Mire Conservation Group and International Peat Society, Finland, 288 pp.
- Joosten, H., Sirin, A., Couwenberg, J., Laine, J. & Smith, P. (in press) The role of peatlands in climate regulation. In: Bonn A., Allott T., Evans, M., Joosten, H. & Stoneman, R. (eds.), *Peatland Restoration and Ecosystem Services - Science, Policy and Practice*. Cambridge University Press, 66–79.
- Joosten, H., Tapio-Biström, M.-L. & Tol, S. (eds.) (2012) *Peatlands - Guidance for Climate Change Mitigation through Conservation, Rehabilitation and Sustainable Use*. Mitigation of Climate Change in Agriculture (MICCA) Programme, FAO/Wetlands International, Rome, 100 pp.
- Jungkunst, H.F. & Fiedler, S. (2007) Latitudinal differentiated water table control of carbon dioxide, methane and nitrous oxide fluxes from hydromorphic soils: feedbacks to climate change. *Global Change Biology*, 13(12), 2668–2683.
- Juottonen, H., Hynninen, A., Nieminen, M., Tuomivirta, T., Tuittila, E.-S., Nousiainen, H., Kell, D.K., Yrjälä, K., Tervahauta, A. & Fritze, H. (2012) Methane-cycling microbial communities and methane emission in natural and restored peatlands. *Applied and Environmental Microbiology*, 78(17), 6386–6389. doi:10.1128/AEM.00261-12.
- Juszczak, R. & Augustin, J. (2013) Exchange of the greenhouse gases methane and nitrous oxide between the atmosphere and a temperate peatland in Central Europe. *Wetlands*, 33(5), 895–907.
- Juutinen, S., Väiliranta, M., Kuutti, V., Laine, A.M., Virtanen, T., Seppä, H., Weckström, J. & Tuittila, E.-S. (2013) Short-term and long-term carbon dynamics in a northern peatland-stream-lake continuum: A catchment approach. *Journal of Geophysical Research: Biogeosciences*, 118, 1–13, doi:10.1002/jgrg.20028.
- Kalacska, M., Arroyo-Mora, J.P., de Gea, J., Snirer, E., Herzog, C. & Moore, T.R. (2013) Videographic analysis of *Eriophorum vaginatum* spatial coverage in an ombrotrophic bog. *Remote Sensing*, 5(12), 6501–6512.
- Kareksela, S., Haapalehto, T., Juutinen, R., Matilainen, R., Tahvanainen, T. & Kotiaho, J.S. (2015) Fighting carbon loss of degraded peatlands by jump-starting ecosystem functioning with ecological restoration. *Science of The Total Environment*, 537, 268–276.
- Kasischke, E.S., Bourgeau-Chavez, L.L., Rober, A.R., Wyatt, K.H., Waddington, J.M. & Turetsky, M.R. (2009) Effects of soil moisture and water depth on ERS SAR backscatter measurements from an Alaskan wetland complex. *Remote Sensing of Environment*, 113(9), 1868–1873.
- Kettridge, N., Turetsky, M.R., Sherwood, J.H., Thompson, D.K., Miller, C.A., Benscoter, B.W., Flannigan, M.D., Wotton, B.M. & Waddington, J.M. (2015) Moderate drop in water table increases peatland vulnerability to post-fire regime shift. *Science Reports*, 5, Article No. 8063, doi:10.1038/srep08063.
- Kivimäki, S.K., Yli-Petäys, M. & Tuittila, E.-S. (2008) Carbon sink function of sedge and *Sphagnum* patches in a restored cut-away peatland: increased functional diversity leads to higher production. *Journal of Applied Ecology*, 45, 921–929.
- Knoth, C., Klein, B., Prinz, T. & Kleinebecker, T. (2013) Unmanned aerial vehicles as innovative remote sensing platforms for high-resolution infrared imagery to support restoration monitoring in cut-over bogs. *Applied Vegetation Science*, 16(3), 509–517.
- Koch, S., Jurasinski, G., Koebsch, F., Koch, M. & Glatzel, S. (2014) Spatial variability of annual estimates of methane emissions in a *Phragmites australis* (Cav.) Trin. ex Steud. dominated restored coastal brackish fen. *Wetlands*, 34(3), 593–602.
- Koebsch, F., Glatzel, S., Hofmann, J., Forbrich, I. &

- Jurasinski, G. (2013a) CO₂ exchange of a temperate fen during the conversion from moderately rewetting to flooding. *Journal of Geophysical Research: Biogeosciences*, 118(2), 940–950.
- Koebisch, F., Glatzel, S. & Jurasinski, G. (2013b) Vegetation controls methane emissions in a coastal brackish fen. *Wetlands Ecology and Management*, 21(5), 323–337.
- Koebisch, F., Jurasinski, G., Koch, M., Hofmann, J. & Glatzel, S. (2015) Controls for multi-scale temporal variation in ecosystem methane exchange during the growing season of a permanently inundated fen. *Agricultural and Forest Meteorology*, 204, 94–105.
- Koehler, A.-K., Murphy, K., Kiely, G. & Sottocornola, M. (2009) Seasonal variation of DOC concentration and annual loss of DOC from an Atlantic blanket bog in South Western Ireland. *Biogeochemistry*, 95(2), 231–242.
- Koehler, A.-K., Sottocornola, M. & Kiely, G. (2011) How strong is the current carbon sequestration of an Atlantic blanket bog? *Global Change Biology*, 17, 309–319.
- Köhl, M., Lasco, R., Cifuentes, M., Jonsson, Ö., Korhonen, K.T., Mundhenk, P., de Jesus Navar, J. & Stinson, G. (2015) Changes in forest production, biomass and carbon: Results from the 2015 UN FAO Global Forest Resource Assessment. *Forest Ecology and Management*, 352, 21–34.
- Kolka, R.K., Grigal, D.F., Verry, E.S. & Nater, E.A. (1999) Mercury and organic carbon relationships in streams draining forested upland peatland watersheds. *Journal of Environmental Quality*, 28, 766–775.
- Komulainen, V.-M., Nykänen, H., Martikainen, P.J. & Laine, J. (1998) Short-term effect of restoration on vegetation change and methane emissions from peatlands drained for forestry in southern Finland. *Canadian Journal of Forest Research*, 28, 402–411.
- Komulainen, V.-M., Tuittila, E.-S., Vasander, H. & Laine, J. (1999) Restoration of drained peatlands in southern Finland: initial effects on vegetation change and CO₂ balance. *Journal of Applied Ecology*, 36, 634–648.
- Koprivnjak, J.F. & Moore, T.R. (1992) Sources, sinks, and fluxes of dissolved organic carbon in subarctic fen catchments. *Arctic and Alpine Research*, 24(3), 204–210.
- Kortelainen, P., Mattsson, T., Finér, L., Ahtiainen, M., Saukkonen, S. & Sallantausta, T. (2006) Controls on the export of C, N, P and Fe from undisturbed boreal catchments, Finland. *Aquatic Sciences*, 68(4), 453–468.
- Kurbatova, J., Li, C., Tataronov, F., Varlagin, A., Shalukhina, N. & Olchev, A. (2009) Modeling of the carbon dioxide fluxes in European Russia peat bogs. *Environmental Research Letters*, 4, 045022, doi:10.1088/1748-9326/4/4/045022.
- Lafleur, P.M., Roulet, N.T. & Admiral, S.W. (2001) Annual cycle of CO₂ exchange at a bog peatland. *Journal of Geophysical Research*, 106(D3), 3071–3081.
- Lähteenoja, O., Reátegui, Y.R., Räsänen, M., Torres, D.D.C., Oinonen, M. & Page, S. (2012) The large Amazonian peatland carbon sink in the subsiding Pastaza-Marañón foreland basin, Peru. *Global Change Biology*, 18(1), 164–178.
- Lähteenoja, O., Ruokolainen, K., Schulman, L. & Oinonen, M. (2009) Amazonian peatlands: an ignored C sink and potential source. *Global Change Biology*, 15(9), 2311–2320.
- Lai, D.Y.F. (2009) Methane dynamics in northern peatlands: a review. *Pedosphere*, 19(4), 409–421.
- Laine, A., Byrne, K.A., Kiely, G. & Tuittila, E.-S. (2007a) Patterns in vegetation and CO₂ dynamics along a water level gradient in a lowland blanket bog. *Ecosystems*, 10, 890–905.
- Laine, A., Wilson, D., Kiely, G. & Byrne, K.A. (2007b) Methane flux dynamics in an Irish lowland blanket bog. *Plant and Soil*, 299(1), 181–193.
- Laine, J., Minkinen, K., Sinisalo, J., Savolainen, I. & Martikainen, P.J. (1997) Greenhouse impact of a mire after drainage for forestry. In: Trettin C.C., Jurgensen M.F., Grigal D.F., Gale, M.R. & Jeglum, J.K. (eds.), *Northern Forested Wetlands, Ecology and Management*. CRC Press, Boca Raton, Florida, 437–447.
- Laine, J., Silvola, J., Tolonen, K., Alm, J., Nykänen, H., Vasander, H., Sallantus, T., Savolainen, I., Sinisalo, J. & Martikainen, P.J. (1996) Effect of water-level drawdown on global warming: Northern peatlands. *Ambio*, 25(3), 179–184.
- Levy, P.E., Burden, A., Cooper, M.D.A., Dinsmore, K.J., Drewer, J., Evans, C., Fowler, D., Gaiawyn, J., Gray, A., Jones, S.K., Jones, T., McNamara, N.P., Mills, R., Ostle, N., Sheppard, L.J., Skiba, U., Sowerby, A., Ward, S.E. & Zielinski, P. (2012) Methane emissions from soils: synthesis and analysis of a large UK data set. *Global Change Biology*, 18(5), 1657–1669.
- Levy, P.E. & Gray, A. (2015) Greenhouse gas balance of a semi-natural peatbog in northern Scotland. *Environmental Research Letters*, 10(9), 094019.
- Li, T., Huang, Y., Zhang, W. & Song, C. (2010)

- CH4MOD_{wetland}: A biogeophysical model for simulating methane emissions from natural wetlands. *Ecological Modelling*, 221(4), 666–680.
- Limpens, J., Berendse, F., Blodau, C., Canadell, J. G., Freeman, C., Holden, J., Roulet, N., Rydin, H. & Schaepman-Strub, G. (2008) Peatlands and the carbon cycle: from local processes to global implications - a synthesis. *Biogeosciences*, 5, 1475–1491.
- Long, K.D., Flanagan, L.B. & Cai, T. (2010) Diurnal and seasonal variation in methane emissions in a northern Canadian peatland measured by eddy covariance. *Global Change Biology*, 16(9), 2420–2435.
- Lund, M., Bjerke, J.W., Drake, B.G., Engelsen, O., Hansen, G.H., Parmentier, F.J.W., Powell, T.L., Silvennoinen, H., Sottocornola, M., Tommervik, H., Weldon, S. & Rasse, D.P. (2015) Low impact of dry conditions on the CO₂ exchange of a Northern-Norwegian blanket bog. *Environmental Research Letters*, 10, 025004.
- Lund, M., Christensen, T.R., Lindroth, A. & Schubert, P. (2012) Effects of drought conditions on the carbon dioxide dynamics in a temperate peatland. *Environmental Research Letters*, 7(4), 045704, 7 pp.
- Lund, M., Lindroth, A., Christensen, T.R. & Strom, L. (2007) Annual CO₂ balance of a temperate bog. *Tellus B*, 59(5), 804–811.
- Maanavilja, L., Riutta, T., Aurela, M., Pulkkinen, M., Laurila, T. & Tuittila, E.-S. (2011) Spatial variation in CO₂ exchange at a northern aapa mire. *Biogeochemistry*, 104, 325–345.
- Maljanen, M., Hytönen, J., Mäkiranta, P., Alm, J., Minkkinen, K., Laine, J. & Martikainen, P.J. (2007) Greenhouse gas emissions from cultivated and abandoned organic croplands in Finland. *Boreal Environment Research*, 12, 133–140.
- Marinier, M., Glatzel, S. & Moore, T.R. (2004) The role of cotton-grass (*Eriophorum vaginatum*) in the exchange of CO₂ and CH₄ at two restored peatlands, eastern Canada. *Ecoscience*, 11(2), 141–149.
- Martikainen, P.J., Nykänen, H., Alm, J. & Silvola, J. (1995) Change in fluxes of carbon dioxide, methane and nitrous oxide due to forest drainage of mire sites of different trophic. *Plant and Soil*, 167–169, 571–577.
- McVeigh, P., Sottocornola, M., Foley, N., Leahy, P. & Kiely, G. (2014) Meteorological and functional response partitioning to explain interannual variability of CO₂ exchange at an Irish Atlantic blanket bog. *Agricultural and Forest Meteorology*, 194, 8–19.
- Melling, L., Goh, K.J., Klioni, A. & Hatano, R. (2012) Is water table the most important factor influencing soil C flux in tropical peatlands? In: *Peatlands in Balance, Proceedings of the 14th International Peat Congress*, International Peat Society, Stockholm, Sweden, Extended Abstract No. 330, 6 pp.
- Melling, L., Hatano, R. & Goh, K.J. (2005) Methane fluxes from three ecosystems in tropical peatland of Sarawak, Malaysia. *Soil Biology and Biochemistry*, 37, 1445–1453.
- Melling, L., Hatano, R. & Goh, K.J. (2007) Nitrous oxide emissions from three ecosystems in tropical peatland of Sarawak, Malaysia. *Soil Science and Plant Nutrition*, 53(6), 792–805.
- Meng, L., Hess, P.G.M., Mahowald, N.M., Yavitt, J.B., Riley, W.J., Subin, Z.M., Lawrence, D.M., Swenson, S.C., Jauhainen, J. & Fuka, D.R. (2012) Sensitivity of wetland methane emissions to model assumptions: application and model testing against site observations. *Biogeosciences*, 9(7), 2793–2819.
- Metzger, C., Jansson, P.E., Lohila, A., Aurela, M., Eickenscheidt, T., Belelli-Marchesini, L., Dinsmore, K., Drewer, J., Van Huissteden, J. & Drösler, M. (2015) CO₂ fluxes and ecosystem dynamics at five European treeless peatlands - merging data and process oriented modeling. *Biogeosciences*, 12, 125–146.
- Minke, M., Augustin, J., Burlo, A., Yarmashuk, T., Chuvashova, H., Thiele, A., Freibauer, A., Tikhonov, V. & Hoffmann, M. (2015) Water level, vegetation composition and plant productivity explain greenhouse gas fluxes in temperate cutover fens after inundation. *Biogeosciences Discussion*, 12(20), 17393–17452.
- Minke, M., Chuvashova, H., Burlo, A., Yarmashuk, T. & Augustin, J. (2012) Measuring GHG emissions from peatlands. In: Tanneberger, F. & Wichtmann, W. (eds.), *Carbon Credits from Peatland Rewetting. Climate - Biodiversity - Land Use*. Schweizerbart, Stuttgart, Germany, 30–36.
- Minkkinen, K. & Laine, J. (2006) Vegetation heterogeneity and ditches create spatial variability in methane fluxes from peatlands drained for forestry. *Plant and Soil*, 285, 289–304.
- Moore, S., Evans, C.D., Page, S.E., Garnett, M.H., Jones, T.G., Freeman, C., Hooijer, A., Wiltshire, A.J., Limin, S.H. & Gauci, V. (2013) Deep instability of deforested tropical peatlands revealed by fluvial organic carbon fluxes.

- Nature*, 493, 660–664.
- Moore, T.R., Matos, L. & Roulet, N.T. (2003) Dynamics and chemistry of dissolved organic carbon in Precambrian Shield catchments and an impounded wetland. *Canadian Journal of Fisheries and Aquatic Sciences*, 60(5), 612–623.
- Müller, D., Warneke, T., Rixen, T., Müller, M., Jamahiri, S., Denis, N., Mujahid, A. & Notholt, J. (2015) Lateral carbon fluxes and CO₂ outgassing from a tropical peat-draining river. *Biogeosciences*, 12(20), 5967–5979.
- Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, L., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T. & Zhang, H. (2013) Anthropogenic and natural radiative forcing. In: Stocker T.F., Qin D., Plattner G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V. & Midgley, P.M. (eds.) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, USA, 659–740.
- Nadeau, D.F., Rousseau, A.N., Coursolle, C., Margolis, H.A. & Parlange, M.B. (2013) Summer methane fluxes from a boreal bog in northern Quebec, Canada, using eddy covariance measurements. *Atmospheric Environment*, 81, 464–474.
- Nagata, O., Takakai, F. & Hatano, R. (2005) Effect of *Sasa* invasion on global warming potential in *Sphagnum* dominated poor fen in Bibai, Japan. *Phyton, Annales Rei Botanicae, Horn*, 45(4), 299–307.
- Nelleman, C. & Corcoran, E. (eds.) (2010) *Dead Planet, Living Planet – Biodiversity and Ecosystem Restoration for Sustainable Development: A Rapid Response Assessment*. GRID-Arendal, Norway for United Nations Environment Programme (UNEP), Nairobi, Kenya, 112 pp.
- Neuzil, S.G. (1997) Onset and rate of peat and carbon accumulation in four domed ombrogenous peat deposits, Indonesia. In: Rieley J.O. & Page S.E. (eds.) *Biodiversity and Sustainability of Tropical Peatlands*, Samara Publishing Ltd., Cardigan, UK, 55–72.
- Nilsson, M., Sagerfors, J., Buffam, I., Laudon, H., Eriksson, T., Grelle, A., Klemetsson, L., Weslien, P.E.R. & Lindroth, A. (2008) Contemporary carbon accumulation in a boreal oligotrophic minerogenic mire – a significant sink after accounting for all C-fluxes. *Global Change Biology*, 14(10), 2317–2332.
- Nykänen, H., Alm, J., Lång, K., Silvola, J. & Martikainen, P.J. (1995) Emissions of CH₄, N₂O and CO₂ from a virgin fen and a fen drained for grassland in Finland. *Journal of Biogeography*, 22, 351–357.
- Nykänen, H., Heikkinen, J.E.P., Pirinen, L., Tiilikainen, K. & Martikainen, P.J. (2003) Annual CO₂ exchange and CH₄ fluxes on a subarctic palsa mire during climatically different years. *Global Biogeochemical Cycles*, 17(1), 1018, doi:10.1029/2002GB001861.
- O'Brien, H.E. (2009) *The Role of Blanket Peat Moorland Management in the Generation and Amelioration of Discolouration of Surface Water Supplies*. PhD thesis, School of Animal, Rural and Environmental Sciences, Nottingham Trent University, 291 pp.
- Olson, D.M., Griffis, T.J., Noormets, A., Kolka, R. & Chen, J. (2013) Interannual, seasonal, and retrospective analysis of the methane and carbon dioxide budgets of a temperate peatland. *Journal of Geophysical Research: Biogeosciences*, 118(1), 226–238.
- Page, S., Hosiolo, A., Wösten, H., Jauhiainen, J., Silvius, M., Rieley, J., Ritzema, H., Tansey, K., Graham, L., Vasander, H. & Limin, S. (2008) Restoration ecology of lowland tropical peatlands in southeast Asia: current knowledge and future research directions. *Ecosystems*, 12(6), 888–905.
- Page, S.E., Rieley, J.O. & Banks, C.J. (2011) Global and regional importance of the tropical peatland carbon pool. *Global Change Biology*, 17(2), 798–818.
- Page, S.E., Siegert, F., Rieley, J.O., Boehm, H.-D.V., Jaya, A. & Limin, S. (2002) The amount of carbon released from peat and forest fires in Indonesia during 1997. *Nature*, 420, 61–65.
- Page, S.E., Wüst, R.A.J., Weiss, D., Rieley, J.O., Shoty, W. & Limin, S.H. (2004) A record of Late Pleistocene and Holocene carbon accumulation and climate change from an equatorial peat bog (Kalimantan, Indonesia): implications for past, present and future carbon dynamics. *Journal of Quaternary Science*, 19(7), 625–635.
- Pangala, S.R., Moore, S., Hornibrook, E.R.C. & Gauci, V. (2013) Trees are major conduits for methane egress from tropical forested wetlands. *New Phytologist*, 197(2), 524–531.
- Peichl, M., Öquist, M., Löfvenius, M.O., Ilstedt, U., Sagerfors, J., Grelle, A., Lindroth, A. & Nilsson, M. (2014) A 12-year record reveals pre-growing

- season temperature and water table level threshold effects on the net carbon dioxide exchange in a boreal fen. *Environmental Research Letters*, 9(5), 055006, 11 pp.
- Petrone, R.M., Waddington, J.M. & Price, J.S. (2003) Ecosystem-scale flux of CO₂ from a restored vacuum harvested peatland. *Wetlands Ecology and Management*, 11(6), 419–432.
- Poopathy, V., Appanah, S. & Durst, P.B. (eds.) (2005) *Helping Forests Take Cover*. RAP Publication 2005/13, Food and Agriculture Organization of the United Nations (FAO), Bangkok, Thailand, 21 pp. Online at: <http://www.fao.org/docrep/008/ae945e/ae945e00.htm>, date accessed 30 Mar 2016.
- Rantakari, M., Mattsson, T., Kortelainen, P., Piirainen, S., Finér, L. & Ahtiainen, M. (2010) Organic and inorganic carbon concentrations and fluxes from managed and unmanaged boreal first-order catchments. *Science of the Total Environment*, 408(7), 1649–1658.
- Regina, K., Nykänen, H., Silvola, J. & Martikainen, P. (1996) Fluxes of nitrous oxide from boreal peatlands as affected by peatland type, water table level and nitrification capacity. *Biogeochemistry*, 35(3), 401–418.
- Renou-Wilson, F., Barry, C., Müller, C. & Wilson, D. (2014) The impacts of drainage, nutrient status and management practice on the full carbon balance of grasslands on organic soils in a maritime temperate zone. *Biogeosciences*, 11(16), 4361–4379.
- Rinne, J., Riutta, T., Pihlatie, M., Aurela, M., Haapanala, S., Tuovinen, J. & Tuittila, E.-S. (2007) Annual cycle of methane emission from a boreal fen measured by the eddy covariance technique. *Tellus*, 59(3) 449–457.
- Riutta, T., Laine, J., Aurela, M., Rinne, J., Vesala, T., Laurila, T., Haapanala, S., Pihlatie, M. & Tuittila, E.-S. (2007) Spatial variation in plant community functions regulates carbon gas dynamics in a boreal fen ecosystem. *Tellus*, 59B(5), 838–852.
- Roehm, C.L. & Roulet, N.T. (2003) Seasonal contribution of CO₂ fluxes in the annual C budget of a northern bog. *Global Biogeochemical Cycles*, 17(1), 29–1–29–9.
- Roulet, N.T., Lafleur, P.M., Richard, P.J.H., Moore, T., Humphreys, E.R. & Bubier, J. (2007) Contemporary carbon balance and late Holocene carbon accumulation in a northern peatland. *Global Change Biology*, 13, 397–411.
- Saarnio, S., Morero, M., Shurpali, N.J., Tuittila, E.-S., Mäkilä, M. & Alm, J. (2007) Annual CO₂ and CH₄ fluxes of pristine boreal mires as a background for the lifecycle analyses of peat energy. *Boreal Environment Research*, 12, 101–113.
- Sagerfors, J., Lindroth, A., Grelle, A., Klemetsson, L., Weslien, P. & Nilsson, M. (2008) Annual CO₂ exchange between a nutrient-poor, minerotrophic, boreal mire and the atmosphere. *Journal of Geophysical Research*, 113(G1), G01001.
- Salm, J.-O., Maddison, M., Tammik, S., Soosaar, K., Truu, J. & Mander, Ü. (2012) Emissions of CO₂, CH₄ and N₂O from undisturbed, drained and mined peatlands in Estonia. *Hydrobiologia*, 692(1), 41–55.
- Samaritani, E., Siegenthaler, A., Yli-Petäys, M., Buttler, A., Christin, P.-A. & Mitchell, E.A.D. (2011) Seasonal net ecosystem carbon exchange of a regenerating cutaway bog: How long does it take to restore the C-sequestration function? *Restoration Ecology*, 19(4), 480–489.
- Schrier-Uijl, A., Veraart, A.J., Leffelaar, P.A., Berendse, F. & Veenendaal, E.M. (2011) Release of CO₂ and CH₄ from lakes and drainage ditches in temperate wetlands. *Biogeochemistry*, 102, 265–279.
- Schulze, E.D., Prokuschkin, A., Arneth, A., Knorre, N. & Vaganov, E.A. (2002) Net ecosystem productivity and peat accumulation in a Siberian aapa mire. *Tellus*, 54B, 531–536.
- Scottish Executive (2007) *ECOSSE - Estimating Carbon in Organic Soils Sequestration and Emissions*. Department of Environment and Rural Affairs, Edinburgh, 165 pp. ISBN: 978 0 7559 1498 2.
- Shannon, R.D. & White, J.R. (1994) A three-year study of controls on methane emissions from two Michigan peatlands. *Biogeochemistry*, 27, 35–60.
- Shannon, R.D., White, J.R., Lawson, J.E. & Gilmour, B.S. (1996) Methane efflux from emergent vegetation in peatlands. *Journal of Ecology*, 84, 239–246.
- Shurpali, N.J., Verma, S.B., Kim, J. & Arkebauer, T.J. (1995) Carbon dioxide exchange in a peatland ecosystem. *Journal of Geophysical Research*, 100(D7), 14319–14326.
- Sirin, A.A., Suvorov, G.G., Chistotin, M.V. & Glagolev, M.V. (2012) Znacheniya emissii metana iz drenaznykh kanav (Values of methane emission from drainage ditches). *Dinamiki okruzhayushchey sredy i global'nyye izmeneniya klimata (Environmental Dynamics and Global Climate Changes)*, 3, 1–10 (in Russian with English abstract). Online at: <http://elibrary.ru/download/97740609.pdf>, accessed 30 Mar 2016.

- Smith, P., Bustamante, M., Ahammad, H., Clark, H., Dong, H., Elsiddig, E.A., Haberl, H., Harper, R., House, J., Jafari, M., Maser, O., Mbow, C., Ravindranath, N.H., Rice, C.W., Robledo Abad, C., Romanovskaya, A., Sperling, F. & Tubiello, F. (2014) Agriculture, Forestry and Other Land Use (AFOLU). In: Edenhofer, O., Pichs-Madruga, R., Sokona, Y., Farahani, E., Kadner, S., Seyboth, K., Adler, A., Baum, I., Brunner, S., Eickemeier, P., Kriemann, B., Savolainen, J., Schlömer, S., von Stechow, C., Zwickel, T. & Minx, J.C. (eds.), *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, USA, 811–922.
- Soegaard, H. & Nordstroem, C. (1999) Carbon dioxide exchange in a high-arctic fen estimated by eddy covariance measurements and modelling. *Global Change Biology*, 5(5), 547–562.
- Soini, P., Riutta, T., Yli-Petäys, M. & Vasander, H. (2010) Comparison of vegetation and CO₂ dynamics between a restored cut-away peatland and a pristine fen: evaluation of the restoration success. *Restoration Ecology*, 18(6), 894–903.
- Sommer, M., Fiedler, S., Glatzel, S. & Kleber, M. (2004) First estimates of regional (Allgäu, Germany) and global CH₄ fluxes from wet colluvial margins of closed depressions in glacial drift areas. *Agriculture, Ecosystems and Environment*, 103(1), 251–257.
- Strack, M., Keith, A.M. & Zu, B. (2014) Growing season carbon dioxide and methane exchange at a restored peatland on the Western Boreal Plain. *Ecological Engineering*, 64, 231–239.
- Strack, M., Waddington, J.A., Bourbonniere, R.A., Buckton, E.L., Shaw, K., Whittington, P. & Price, J.S. (2008) Effect of water table drawdown on peatland dissolved organic carbon export and dynamics. *Hydrological Processes*, 22(17), 3373–3385, doi:10.1002/hyp.6931.
- Strack, M., Waddington, J.M., Rochefort, L. & Tuittila, E.-S. (2006) Response of vegetation and net ecosystem carbon dioxide exchange at different peatland microforms following water table drawdown. *Journal of Geophysical Research*, 111(G2), doi:10.1029/2005JG000145.
- Strack, M. & Zuback, Y.C.A. (2013) Annual carbon balance of a peatland 10 yr following restoration. *Biogeosciences*, 10, 2885–2896.
- Suyker, A.E., Verma, S.B. & Arkebauer, T.J. (1997) Season-long measurement of carbon dioxide exchange in a boreal fen. *Journal of Geophysical Research*, 102(D24), 29021–29028.
- Tauchnitz, N., Brumme, R., Bernsdorf, S. & Meissner, R. (2008) Nitrous oxide and methane fluxes of a pristine slope mire in the German National Park Harz Mountains. *Plant and Soil*, 303(1–2), 131–138.
- Torbick, N., Persson, A., Olefeldt, D., Froelking, S., Salas, W., Hagen, S., Crill, P. & Li, C. (2012) High resolution mapping of peatland hydroperiod at a high-latitude Swedish mire. *Remote Sensing*, 4(7), 1974–1994.
- Tuittila, E.-S., Komulainen, V.-M., Vasander, H. & Laine, J. (1999) Restored cut-away peatland as a sink for atmospheric CO₂. *Oecologia*, 120, 563–574.
- Tuittila, E.-S., Komulainen, V.-M., Vasander, H., Nykänen, H., Martikainen, P.J. & Laine, J. (2000) Methane dynamics of a restored cut-away peatland. *Global Change Biology*, 6, 569–581.
- Turetsky, M.R., Benscoter, B., Page, S., Rein, G., van der Werf, G.R. & Watts, A. (2015) Global vulnerability of peatlands to fire and carbon loss. *Nature Geoscience*, 8(1), 11–14.
- Turner, E.K., Worrall, F. & Burt, T.P. (2013) The effect of drain blocking on the dissolved organic carbon (DOC) budget of an upland peat catchment in the UK. *Journal of Hydrology*, 479, 169–179.
- Urban, N.R., Bayley, S.E. & Eisenreich, S.J. (1989) Export of dissolved organic carbon and acidity from peatlands. *Water Resources Research*, 25(7), 1619–1628.
- Urbanová, Z. (2012) Vegetation and carbon gas dynamics under a changed hydrological regime in central European peatlands. *Plant Ecology and Diversity*, 5(1), 89–103.
- Urbanová, Z., Bárta, J. & Pícek, T. (2013a) Methane emissions and methanogenic Archaea on pristine, drained and restored mountain peatlands, central Europe. *Ecosystems*, 16(4), 664–677.
- Urbanová, Z., Pícek, T. & Tuittila, E.-S. (2013b) Sensitivity of carbon gas fluxes to weather variability on pristine, drained and rewetted temperate bogs. *Mires and Peat*, 11(04), 1–14.
- van den Pol-van Dasselaar, A., Corré, W.J., Priemé, A., Klemedtsson, Å.K., Weslien, P., Klemedtsson, L., Stein, A. & Oenema, O. (1998) Spatial variability of methane, nitrous oxide, and carbon dioxide emissions from drained grasslands. *Soil Science Society of America Journal*, 62(3), 810–817.
- Vanselow-Algan, M., Schmidt, S.R., Greven, M., Fiencke, C., Kutzbach, L. & Pfeiffer, E.M. (2015) High methane emissions dominated

- annual greenhouse gas balances 30 years after bog rewetting. *Biogeosciences*, 12(14), 4361–4371.
- van Winden, J.F., Reichart, G.J., McNamara, N.P., Benthien, A. & Damsté, J.S.S. (2012) Temperature-induced increase in methane release from peat bogs: a mesocosm experiment. *PLoS ONE*, 7(6), e39614. doi:10.1371/journal.pone.0039614.
- Verma, S.B., Ullman, F.G., Billesbach, D., Clement, R.J., Kim, J. & Verry, E.S. (1992) Eddy-correlation measurements of methane flux in a northern peatland ecosystem. *Boundary-Layer Meteorology*, 58(3), 289–304.
- von Arnold, K. (2004) *Forests and Greenhouse Gases: Fluxes of CO₂, CH₄ and N₂O from Drained Forests on Organic Soils*. Thesis, Linköping Studies in Arts and Science No. 302, Linköping University, Sweden, 47 pp. Online at: <https://www.diva-portal.org/smash/get/diva2:20781/FULLTEXT01.pdf>, accessed 23 Mar 2016.
- Waddington, J.M. & Day, S.M. (2007) Methane emissions from a peatland following restoration. *Journal of Geophysical Research*, 112(G3), G03018.
- Waddington, J.M. & Price, J.S. (2000) Effect of peatland drainage, harvesting and restoration on atmospheric water and carbon exchange. *Physical Geography*, 21(5), 433–451.
- Waddington, J.M. & Roulet, N.T. (2000) Carbon balance of a boreal patterned peatland. *Global Change Biology*, 6(1), 87–97.
- Waddington, J.M., Strack, M. & Greenwood, M.J. (2010) Toward restoring the net carbon sink function of degraded peatlands: Short-term response in CO₂ exchange to ecosystem-scale restoration. *Journal of Geophysical Research*, 115, G01008, doi:10.1029/2009JG001090.
- Waddington, J.M., Tóth, K. & Bourbonniere, R.A. (2008) Dissolved organic carbon export from a cutover and restored peatland. *Hydrological Processes*, 22, 2215–2224.
- Walter, B.P., Heimann, M. & Matthews, E. (2001) Modeling modern methane emissions from natural wetlands 1. Model description and results. *Journal of Geophysical Research*, 106(D24), 34189–34206.
- Whiting, G.J. & Chanton, J.P. (2001) Greenhouse carbon balance of wetlands: methane emission versus carbon sequestration. *Tellus*, 53B, 521–528.
- Wickland, K. (2001) Carbon gas exchange at a southern Rocky Mountain wetland, 1996–1998. *Global Biogeochemical Cycles*, 15(2), 321–335.
- Wild, U., Kamp, T., Lenz, A., Heinz, S. & Pfadenhauer, J. (2001) Cultivation of *Typha* spp. in constructed wetlands for peatland restoration. *Ecological Engineering*, 17, 49–54.
- Wilson, D., Alm, J., Laine, J., Byrne, K.A., Farrell, E.P. & Tuittila, E.-S. (2009) Rewetting of cutaway peatlands: Are we re-creating hotspots of methane emissions? *Restoration Ecology*, 17(6), 796–806.
- Wilson, D., Farrell, C.A., Müller, C., Hepp, S. & Renou-Wilson, F. (2013) Rewetted industrial cutaway peatlands in western Ireland: prime location for climate change mitigation? *Mires and Peat*, 11(01), 1–22.
- Wilson, D., Tuittila, E.-S., Alm, J., Laine, J., Farrell, E.P. & Byrne, K.A. (2007) Carbon dioxide dynamics of a restored maritime peatland. *Ecoscience*, 14(1), 71–80.
- Yli-Petäys, M., Laine, J., Vasander, H. & Tuittila, E.-S. (2007) Carbon gas exchange of a re-vegetated cut-away peatland five decades after abandonment. *Boreal Environment Research*, 12, 177–190.
- Yu, Z.C. (2012) Northern peatland carbon stocks and dynamics: a review. *Biogeosciences*, 9, 4071–4085.
- Yule, C. & Gomez, L. (2009) Leaf litter decomposition in a tropical peat swamp forest in Peninsular Malaysia. *Wetlands Ecology and Management*, 17(3), 231–241.
- Zulkifli, Y. (2002) Hydrological attributes of a disturbed peat swamp forest. In: Parish, F., Padmanabhan, E., Lee, C.L. & Thang, H.C. (eds.), *Prevention and Control of Fire in Peatlands*, Proceedings of Workshop at Forestry Training Unit, Kepong, Kuala Lumpur, 19–21 March 2002, Forestry Department Peninsular Malaysia, Kuala Lumpur and Global Environment Centre, Selangor, 51–55.

Submitted 20 Jan 2016, final revision 14 Mar 2016
 Editor: Olivia Bragg

Author for correspondence:

Dr David Wilson, Earthy Matters Environmental Consultants, Glenvar, Co. Donegal, Ireland.
 Tel: + 353 74 9177613; E-mail: david.wilson@earthymatters.ie